

## HAPLOTYPES OF THE TNFRSF11B GENE

### RELATED APPLICATIONS

This application is a continuation-in-part of International Application PCT/US00/18803 filed July 10, 2000, which claims the benefit of U.S. Provisional Application Serial No. 60/143,020 filed July 9, 1999.

### FIELD OF THE INVENTION

This invention relates to variation in genes that encode pharmaceutically-important proteins. In particular, this invention provides genetic variants of the human tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin) (TNFRSF11B) gene and methods for identifying which variant(s) of this gene is/are possessed by an individual.

### BACKGROUND OF THE INVENTION

Current methods for identifying pharmaceuticals to treat disease often start by identifying, cloning, and expressing an important target protein related to the disease. A determination of whether an agonist or antagonist is needed to produce an effect that may benefit a patient with the disease is then made. Then, vast numbers of compounds are screened against the target protein to find new potential drugs. The desired outcome of this process is a lead compound that is specific for the target, thereby reducing the incidence of the undesired side effects usually caused by activity at non-intended targets. The lead compound identified in this screening process then undergoes further *in vitro* and *in vivo* testing to determine its absorption, disposition, metabolism and toxicological profiles. Typically, this testing involves use of cell lines and animal models with limited, if any, genetic diversity.

What this approach fails to consider, however, is that natural genetic variability exists between individuals in any and every population with respect to pharmaceutically-important proteins, including the protein targets of candidate drugs, the enzymes that metabolize these drugs and the proteins whose activity is modulated by such drug targets. Subtle alteration(s) in the primary nucleotide sequence of a gene encoding a pharmaceutically-important protein may be manifested as significant variation in expression, structure and/or function of the protein. Such alterations may explain the relatively high degree of uncertainty inherent in the treatment of individuals with a drug whose design is based upon a single representative example of the target or enzyme(s) involved in metabolizing the drug. For example, it is well-established that some drugs frequently have lower efficacy in some individuals than others, which means such individuals and their physicians must weigh the possible benefit of a larger dosage against a

greater risk of side effects. Also, there is significant variation in how well people metabolize drugs and other exogenous chemicals, resulting in substantial interindividual variation in the toxicity and/or efficacy of such exogenous substances (Evans et al., 1999, *Science* 286:487-491). This variability in efficacy or toxicity of a drug in genetically-diverse patients makes many drugs ineffective or even dangerous in certain groups of the population, leading to the failure of such drugs in clinical trials or their early withdrawal from the market even though they could be highly beneficial for other groups in the population. This problem significantly increases the time and cost of drug discovery and development, which is a matter of great public concern.

It is well-recognized by pharmaceutical scientists that considering the impact of the genetic variability of pharmaceutically-important proteins in the early phases of drug discovery and development is likely to reduce the failure rate of candidate and approved drugs (Marshall A 1997 *Nature Biotech* 15:1249-52; Kleyn PW et al. 1998 *Science* 281: 1820-21; Kola I 1999 *Curr Opin Biotech* 10:589-92; Hill AVS et al. 1999 in *Evolution in Health and Disease* Stearns SS (Ed.) Oxford University Press, New York, pp 62-76; Meyer U.A. 1999 in *Evolution in Health and Disease* Stearns SS (Ed.) Oxford University Press, New York, pp 41-49; Kalow W et al. 1999 *Clin. Pharm. Therap.* 66:445-7; Marshall, E 1999 *Science* 284:406-7; Judson R et al. 2000 *Pharmacogenomics* 1:1-12; Roses AD 2000 *Nature* 405:857-65). However, in practice this has been difficult to do, in large part because of the time and cost required for discovering the amount of genetic variation that exists in the population (Chakravarti A 1998 *Nature Genet* 19:216-7; Wang DG et al 1998 *Science* 280:1077-82; Chakravarti A 1999 *Nat Genet* 21:56-60 (suppl); Stephens JC 1999 *Mol. Diagnosis* 4:309-317; Kwok PY and Gu S 1999 *Mol. Med. Today* 5:538-43; Davidson S 2000 *Nature Biotech* 18:1134-5).

The standard for measuring genetic variation among individuals is the haplotype, which is the ordered combination of polymorphisms in the sequence of each form of a gene that exists in the population. Because haplotypes represent the variation across each form of a gene, they provide a more accurate and reliable measurement of genetic variation than individual polymorphisms. For example, while specific variations in gene sequences have been associated with a particular phenotype such as disease susceptibility (Roses AD *supra*; Ulbrecht M et al. 2000 *Am J Respir Crit Care Med* 161: 469-74) and drug response (Wolfe CR et al. 2000 *BMJ* 320:987-90; Dahl BS 1997 *Acta Psychiatr Scand* 96 (Suppl 391): 14-21), in many other cases an individual polymorphism may be found in a variety of genomic backgrounds, i.e., different haplotypes, and therefore shows no definitive coupling between the polymorphism and the causative site for the phenotype (Clark AG et al. 1998 *Am J Hum Genet* 63:595-612; Ulbrecht M et al. 2000 *supra*; Drysdale et al. 2000 *PNAS* 97:10483-10488). Thus, there is an unmet need in the pharmaceutical industry for information on what haplotypes exist in the population for pharmaceutically-important genes. Such haplotype information would be useful in improving the

efficiency and output of several steps in the drug discovery and development process, including target validation, identifying lead compounds, and early phase clinical trials (Marshall et al., *supra*).

One pharmaceutically-important gene for the treatment of osteopetrosis, osteoporosis, metastatic bone disease, Paget's disease, rheumatoid arthritis, and periodontal bone disease is the tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin) (TNFRSF11B) gene or its encoded product. TNFRSF11B, also known as osteoclastogenesis inhibitory factor (OCIF or OIF) or Osteoprotegerin (OPG) (Simonet et al., 90 *Cell* 89(2):309-319, 1997), is a soluble member of the tumor necrosis factor receptor (TNFR) superfamily, and binds to at least one TNF-related cytokine, RANKL (also known as TRANCE, OPGL, ODF), which stimulates differentiation of osteoclasts, which are the bone resorbing cells in the body.

*In vitro* studies have shown that TNFRSF11B neutralizes RANKL-induced osteoclastogenesis by binding to RANKL, suggesting that TNFRSF11B is actually a secreted "decoy" receptor for RANKL that blocks initiation of a critical RANK-RANKL signal transduction pathway within osteoclast precursor cells (Takahashi et al., *Biochem. Biophys. Res. Commun.* 256:449-455, 1999). As a result of this blocking action, the number of mature osteoclasts is decreased. *In vivo*, TNFRSF11B increases bone mineral density and bone volume in normal rats and also exhibits hypocalcemic effects in normal mice and in hypercalcemic nude mice carrying tumors associated with humoral hypercalcemia of malignancy (Akatsu et al., *Bone* 23:495-498, 1998). Also, it was reported that TNFRSF11B knock-out mice develop severe osteoporosis due to enhanced osteoclastogenesis when they grew to be adults (Mizuno et al., *Biochem. Biophys. Res. Commun.* 247:610-615, 1998). Thus, along with RANKL and RANK, TNFRSF11B is one of the key molecules that regulate osteoclast recruitment and function, and as such, an understanding of variation in the TNFRSF11B gene should be useful in developing new therapies for metabolic diseases caused by abnormal osteoclast recruitment and function such as osteopetrosis, osteoporosis, metastatic bone disease, Paget's disease, rheumatoid arthritis, and periodontal bone disease.

The tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin) gene is located on chromosome 8q24 and contains 5 exons that encode a 401 amino acid protein. A reference sequence for the TNFRSF11B gene is shown in the contiguous lines of Figure 1 (Genaissance Reference No. 3826804; SEQ ID NO: 1). Reference sequences for the coding sequence (GenBank Accession No. GPI\_19442.1) and protein are shown in Figures 2 (SEQ ID NO: 2) and 3 (SEQ ID NO: 3), respectively.

Within the TNFR superfamily, TNFRSF11B is most similar to TNFRII and CD40, in that this secreted protein has no transmembrane segment, and circulates as a disulfide-linked homodimer. TNFRSF11B has four cysteine-rich domains and two death domain homologous regions present in tandem at the C-terminal portion of the protein (Morinaga et al., *Eur. J. Biochem.* 254:6850691, 1998; Mizuno et al., *Gene* 214:339-343, 1998).

One group has reported finding single nucleotide polymorphisms (SNPs) at positions 9 and 22 of exon 1, which correspond to positions 1009 and 1022 in Figure 1 (Yasuda et al., *Endocrinology* 139:1329-1337, 1997; GENBANK Acc. No. E15271.1). The SNPs at these sites would result in variation in the encoded amino acid sequence at position 3 (lysine or asparagine) and/or position 8 (alanine or serine), respectively, depending on the particular combination of nucleotides found at these polymorphic sites in one of the two copies of the TNFRSF11B gene from an individual. The variation of a guanine or cytosine at position 1009 in Figure 1 is referred to herein as PS5.

Because of the potential for variation in the TNFRSF11B gene to affect the expression and function of the encoded protein, it would be useful to know whether additional polymorphisms exist in the TNFRSF11B gene, as well as how such polymorphisms are combined in different copies of the gene. Such information could be applied for studying the biological function of TNFRSF11B as well as in identifying drugs targeting this protein for the treatment of disorders related to its abnormal expression or function.

#### SUMMARY OF THE INVENTION

Accordingly, the inventors herein have discovered 18 novel polymorphic sites in the TNFRSF11B gene. These polymorphic sites (PS) correspond to the following nucleotide positions in Figure 1: 504 (PS1), 717 (PS2), 744 (PS3), 778 (PS4), 1045 (PS6), 1122 (PS7), 1218 (PS8), 2014 (PS9), 2177 (PS10), 5906 (PS11), 6010 (PS12), 8110 (PS13), 8333 (PS14), 8354 (PS15), 8402 (PS16), 8459 (PS17), 10203 (PS18) and 10512 (PS19). The polymorphisms at these sites are guanine or thymine at PS1, cytosine or thymine at PS2, guanine or thymine at PS3, thymine or cytosine at PS4, cytosine or thymine at PS6, guanine or adenine at PS7, cytosine or adenine at PS8, cytosine or thymine at PS9, thymine or cytosine at PS10, thymine or cytosine at PS11, cytosine or thymine at PS12, guanine or adenine at PS13, cytosine or thymine at PS14, adenine or guanine at PS15, adenine or guanine at PS16, adenine or cytosine at PS17, guanine or adenine at PS18 and thymine or cytosine at PS19. In addition, the inventors have determined the identity of the alleles at these sites, as well as at the previously identified site at nucleotide position 1009 (PS5), in a human reference population of 79 unrelated individuals self-identified as belonging to one of four major population groups: African descent, Asian, Caucasian and Hispanic/Latino. From this information, the inventors deduced a set of haplotypes and haplotype pairs for PS1-PS19 in the TNFRSF11B gene, which are shown below in Tables 5 and 4, respectively. Each of these TNFRSF11B haplotypes constitutes a code that defines the variant nucleotides that exist in the human population at this set of polymorphic sites in the TNFRSF11B gene. Thus each TNFRSF11B haplotype also represents a naturally-occurring isoform (also referred to herein as an "isogene") of the TNFRSF11B gene. The frequency of each haplotype and haplotype pair within the total reference population and within each of the

four major population groups included in the reference population was also determined.

Thus, in one embodiment, the invention provides a method, composition and kit for genotyping the TNFRSF11B gene in an individual. The genotyping method comprises identifying the nucleotide pair that is present at one or more polymorphic sites selected from the group consisting of PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19 in both copies of the TNFRSF11B gene from the individual. A genotyping composition of the invention comprises an oligonucleotide probe or primer which is designed to specifically hybridize to a target region containing, or adjacent to, one of these novel TNFRSF11B polymorphic sites. A genotyping kit of the invention comprises a set of oligonucleotides designed to genotype each of these novel TNFRSF11B polymorphic sites. In a preferred embodiment, the genotyping kit comprises a set of oligonucleotides designed to genotype each of PS1-PS19. The genotyping method, composition, and kit are useful in determining whether an individual has one of the haplotypes in Table 5 below or has one of the haplotype pairs in Table 4 below.

The invention also provides a method for haplotyping the TNFRSF11B gene in an individual. In one embodiment, the haplotyping method comprises determining, for one copy of the TNFRSF11B gene, the identity of the nucleotide at one or more polymorphic sites selected from the group consisting of PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19. In another embodiment, the haplotyping method comprises determining whether one copy of the individual's TNFRSF11B gene is defined by one of the TNFRSF11B haplotypes shown in Table 5, below, or a sub-haplotype thereof. In a preferred embodiment, the haplotyping method comprises determining whether both copies of the individual's TNFRSF11B gene are defined by one of the TNFRSF11B haplotype pairs shown in Table 4 below, or a sub-haplotype pair thereof. Establishing the TNFRSF11B haplotype or haplotype pair of an individual is useful for improving the efficiency and reliability of several steps in the discovery and development of drugs for treating diseases associated with TNFRSF11B activity, e.g., osteopetrosis, osteoporosis, metastatic bone disease, Paget's disease, rheumatoid arthritis, and periodontal bone disease.

For example, the haplotyping method can be used by the pharmaceutical research scientist to validate TNFRSF11B as a candidate target for treating a specific condition or disease, such as osteoporosis, predicted to be associated with TNFRSF11B activity. Determining for a particular population the frequency of one or more of the individual TNFRSF11B haplotypes or haplotype pairs described herein will facilitate a decision on whether to pursue TNFRSF11B as a target for treating the specific disease of interest. In particular, if variable TNFRSF11B activity is associated with the disease, then one or more TNFRSF11B haplotypes or haplotype pairs will be found at a higher frequency in disease cohorts than in appropriately genetically matched controls. Conversely, if each of the observed

10044674-010902  
5 TNFRSF11B haplotypes are of similar frequencies in the disease and control groups, then it may be inferred that variable TNFRSF11B activity has little, if any, involvement with that disease. In either case, the pharmaceutical research scientist can, without *a priori* knowledge as to the phenotypic effect of any TNFRSF11B haplotype or haplotype pair, apply the information derived from detecting TNFRSF11B haplotypes in an individual to decide whether modulating TNFRSF11B activity would be useful in treating the disease.

10 The claimed invention is also useful in screening for compounds targeting TNFRSF11B to treat a specific condition or disease, such as osteoporosis, predicted to be associated with TNFRSF11B activity. For example, detecting which of the TNFRSF11B haplotypes or haplotype pairs disclosed herein are present in individual members of a population with the specific disease of interest enables the pharmaceutical scientist to screen for a compound(s) that displays the highest desired agonist or antagonist activity for each of the TNFRSF11B isoforms present in the disease population, or for only the most frequent TNFRSF11B isoforms present in the disease population. Thus, without requiring any *a priori* knowledge of the phenotypic effect of any particular TNFRSF11B haplotype or haplotype pair, the claimed haplotyping method provides the scientist with a tool to identify lead compounds that are more likely to show efficacy in clinical trials.

20 Haplotyping the TNFRSF11B gene in an individual is also useful in the design of clinical trials of candidate drugs for treating a specific condition or disease, such as osteoporosis, predicted to be associated with TNFRSF11B activity. For example, instead of randomly assigning patients with the disease of interest to the treatment or control group as is typically done now, determining which of the TNFRSF11B haplotype(s) disclosed herein are present in individual patients enables the pharmaceutical scientist to distribute TNFRSF11B haplotypes and/or haplotype pairs evenly to treatment and control groups, thereby reducing the potential for bias in the results that could be introduced by a larger frequency of a TNFRSF11B haplotype or haplotype pair that is associated with response to the drug being studied in the trial, even if this association was previously unknown. Thus, by practicing the claimed invention, the scientist can more confidently rely on the information learned from the trial, without first determining the phenotypic effect of any TNFRSF11B haplotype or haplotype pair.

30 In another embodiment, the invention provides a method for identifying an association between a trait and a TNFRSF11B genotype, haplotype, or haplotype pair for one or more of the novel polymorphic sites described herein. The method comprises comparing the frequency of the TNFRSF11B genotype, haplotype, or haplotype pair in a population exhibiting the trait with the frequency of the TNFRSF11B genotype or haplotype in a reference population. A higher frequency of the TNFRSF11B genotype, haplotype, or haplotype pair in the trait population than in the reference population indicates the trait is associated with the TNFRSF11B genotype, haplotype, or haplotype pair. In preferred embodiments, the

trait is susceptibility to a disease, severity of a disease, the staging of a disease or response to a drug. In a particularly preferred embodiment, the TNFRSF11B haplotype is selected from the haplotypes shown in Table 5, or a sub-haplotype thereof. Such methods have applicability in developing diagnostic tests and therapeutic treatments for osteopetrosis, osteoporosis, metastatic bone disease, Paget's disease, rheumatoid arthritis, and periodontal bone disease.

In yet another embodiment, the invention provides an isolated polynucleotide comprising a nucleotide sequence which is a polymorphic variant of a reference sequence for the TNFRSF11B gene or a fragment thereof. The reference sequence comprises the contiguous sequences shown in Figure 1 and the polymorphic variant comprises at least one polymorphism selected from the group consisting of thymine at PS1, thymine at PS2, thymine at PS3, cytosine at PS4, thymine at PS6, adenine at PS7, adenine at PS8, thymine at PS9, cytosine at PS10, cytosine at PS11, thymine at PS12, adenine at PS13, thymine at PS14, guanine at PS15, guanine at PS16, cytosine at PS17, adenine at PS18 and cytosine at PS19. In a preferred embodiment, the polymorphic variant comprises an additional polymorphism of cytosine at PS5.

A particularly preferred polymorphic variant is an isogene of the TNFRSF11B gene. A TNFRSF11B isogene of the invention comprises guanine or thymine at PS1, cytosine or thymine at PS2, guanine or thymine at PS3, thymine or cytosine at PS4, guanine or cytosine at PS5, cytosine or thymine at PS6, guanine or adenine at PS7, cytosine or adenine at PS8, cytosine or thymine at PS9, thymine or cytosine at PS10, thymine or cytosine at PS11, cytosine or thymine at PS12, guanine or adenine at PS13, cytosine or thymine at PS14, adenine or guanine at PS15, adenine or guanine at PS16, adenine or cytosine at PS17, guanine or adenine at PS18 and thymine or cytosine at PS19. The invention also provides a collection of TNFRSF11B isogenes, referred to herein as a TNFRSF11B genome anthology.

In another embodiment, the invention provides a polynucleotide comprising a polymorphic variant of a reference sequence for a TNFRSF11B cDNA or a fragment thereof. The reference sequence comprises SEQ ID NO:2 (Fig.2) and the polymorphic cDNA comprises at least one polymorphism selected from the group consisting of thymine at a position corresponding to nucleotide 699, guanine at a position corresponding to nucleotide 720, guanine at a position corresponding to nucleotide 768, adenine at a position corresponding to nucleotide 841 and cytosine at a position corresponding to nucleotide 1150. In a preferred embodiment, the polymorphic variant comprises an additional polymorphism of cytosine at a position corresponding to nucleotide 9. A particularly preferred polymorphic cDNA variant comprises the coding sequence of a TNFRSF11B isogene defined by haplotypes 1-3, 7-14, 18, 21 and 22.

Polynucleotides complementary to these TNFRSF11B genomic and cDNA variants are also provided by the invention. It is believed that polymorphic variants of the TNFRSF11B gene will be useful in studying the expression and function of TNFRSF11B, and in expressing TNFRSF11B protein for use in screening for candidate drugs to treat diseases related to TNFRSF11B activity.

In other embodiments, the invention provides a recombinant expression vector comprising one of the polymorphic genomic and cDNA variants operably linked to expression regulatory elements as well as a recombinant host cell transformed or transfected with the expression vector. The recombinant vector and host cell may be used to express TNFRSF11B for protein structure analysis and drug binding studies.

In yet another embodiment, the invention provides a polypeptide comprising a polymorphic variant of a reference amino acid sequence for the TNFRSF11B protein. The reference amino acid sequence comprises SEQ ID NO:3 (Fig.3) and the polymorphic variant comprises at least one variant amino acid selected from the group consisting of methionine at a position corresponding to amino acid position 240 and methionine at a position corresponding to amino acid position 281. In some embodiments, the polymorphic variant also comprises asparagine at a position corresponding to amino acid position 3. A polymorphic variant of TNFRSF11B is useful in studying the effect of the variation on the biological activity of TNFRSF11B as well as on the binding affinity of candidate drugs targeting TNFRSF11B for the treatment of osteopetrosis, osteoporosis, metastatic bone disease, Paget's disease, rheumatoid arthritis, and periodontal bone disease.

The present invention also provides antibodies that recognize and bind to the above polymorphic TNFRSF11B protein variant. Such antibodies can be utilized in a variety of diagnostic and prognostic formats and therapeutic methods.

The present invention also provides nonhuman transgenic animals comprising one or more of the TNFRSF11B polymorphic genomic variants described herein and methods for producing such animals. The transgenic animals are useful for studying expression of the TNFRSF11B isogenes *in vivo*, for *in vivo* screening and testing of drugs targeted against TNFRSF11B protein, and for testing the efficacy of therapeutic agents and compounds for osteopetrosis, osteoporosis, metastatic bone disease, Paget's disease, rheumatoid arthritis, and periodontal bone disease in a biological system.

The present invention also provides a computer system for storing and displaying polymorphism data determined for the TNFRSF11B gene. The computer system comprises a computer processing unit; a display; and a database containing the polymorphism data. The polymorphism data includes one or more of the following: the polymorphisms, the genotypes, the haplotypes, and the haplotype pairs identified for the TNFRSF11B gene in a reference population. In a preferred embodiment, the computer system is capable of producing a display showing TNFRSF11B haplotypes organized according to their evolutionary relationships.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a reference sequence for the TNFRSF11B gene (Genaissance Reference No. 3826804; contiguous lines), with the start and stop positions of each region of coding sequence indicated

with a bracket ([ or ]) and the numerical position below the sequence and the polymorphic site(s) and polymorphism(s) identified by Applicants in a reference population indicated by the variant nucleotide positioned below the polymorphic site in the sequence. SEQ ID NO:1 is equivalent to Figure 1, with the two alternative allelic variants of each polymorphic site indicated by the appropriate nucleotide symbol (R= G or A, Y= T or C, M= A or C, K= G or T, S= G or C, and W= A or T; WIPO standard ST.25). SEQ ID NO:94 is a modified version of SEQ ID NO:1 that shows the context sequence of each polymorphic site, PS1-PS19, in a uniform format to facilitate electronic searching. For each polymorphic site, SEQ ID NO:94 contains a block of 60 bases of the nucleotide sequence encompassing the centrally-located polymorphic site at the 30<sup>th</sup> position, followed by 60 bases of unspecified sequence to represent that each PS is separated by genomic sequence whose composition is defined elsewhere herein.

Figure 2 illustrates a reference sequence for the TNFRSF11B coding sequence (contiguous lines; SEQ ID NO:2), with the polymorphic site(s) and polymorphism(s) identified by Applicants in a reference population indicated by the variant nucleotide positioned below the polymorphic site in the sequence.

Figure 3 illustrates a reference sequence for the TNFRSF11B protein (contiguous lines; SEQ ID NO:3), with the variant amino acid(s) caused by the polymorphism(s) of Figure 2 positioned below the polymorphic site in the sequence.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is based on the discovery of novel variants of the TNFRSF11B gene. As described in more detail below, the inventors herein discovered 22 isogenes of the TNFRSF11B gene by characterizing the TNFRSF11B gene found in genomic DNAs isolated from an Index Repository that contains immortalized cell lines from one chimpanzee and 93 human individuals. The human individuals included a reference population of 79 unrelated individuals self-identified as belonging to one of four major population groups: Caucasian (21 individuals), African descent (20 individuals), Asian (20 individuals), or Hispanic/Latino (18 individuals). To the extent possible, the members of this reference population were organized into population subgroups by their self-identified ethnogeographic origin as shown in Table 1 below. In addition, the Index Repository contains three unrelated indigenous American Indians (one from each of North, Central and South America), one three-generation Caucasian family (from the CEPH Utah cohort) and one two-generation African-American family.

Table 1. Population Groups in the Index Repository

Population Group	Population Subgroup	No. of Individuals
African descent		20
	Sierra Leone	1
Asian		20
	Burma	1
	China	3
	Japan	6
	Korea	1
	Philippines	5
	Vietnam	4
Caucasian		21
	British Isles	3
	British Isles/Central	4
	British Isles/Eastern	1
	Central/Eastern	1
	Eastern	3
	Central/Mediterranean	1
	Mediterranean	2
	Scandinavian	2
Hispanic/Latino		18
	Caribbean	8
	Caribbean (Spanish Descent)	2
	Central American (Spanish Descent)	1
	Mexican American	4
	South American (Spanish Descent)	3

The TNFRSF11B isogenes present in the human reference population are defined by haplotypes  
 5 for 19 polymorphic sites in the TNFRSF11B gene, 18 of which are believed to be novel. The  
 TNFRSF11B polymorphic sites identified by the inventors are referred to as PS1-PS19 to designate the  
 order in which they are located in the gene (see Table 3 below), with the novel polymorphic sites referred  
 to as PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18  
 and PS19. Using the genotypes identified in the Index Repository for PS1-PS19 and the methodology  
 10 described in the Examples below, the inventors herein also determined the pair of haplotypes for the  
 TNFRSF11B gene present in individual human members of this repository. The human genotypes and  
 haplotypes found in the repository for the TNFRSF11B gene include those shown in Tables 4 and 5,  
 respectively. The polymorphism and haplotype data disclosed herein are useful for validating whether  
 TNFRSF11B is a suitable target for drugs to treat osteopetrosis, osteoporosis, metastatic bone disease,  
 15 Paget's disease, rheumatoid arthritis, and periodontal bone disease, screening for such drugs and reducing  
 bias in clinical trials of such drugs.

In the context of this disclosure, the following terms shall be defined as follows unless otherwise

indicated:

**Allele** - A particular form of a genetic locus, distinguished from other forms by its particular nucleotide sequence.

**Candidate Gene** - A gene which is hypothesized to be responsible for a disease, condition, or the response to a treatment, or to be correlated with one of these.

**Gene** - A segment of DNA that contains the coding sequence for a protein, wherein the segment may include promoters, exons, introns, and other untranslated regions that control expression.

**Genotype** - An unphased 5' to 3' sequence of nucleotide pair(s) found at one or more polymorphic sites in a locus on a pair of homologous chromosomes in an individual. As used herein, genotype includes a full-genotype and/or a sub-genotype as described below.

**Full-genotype** - The unphased 5' to 3' sequence of nucleotide pairs found at all polymorphic sites examined herein in a locus on a pair of homologous chromosomes in a single individual.

**Sub-genotype** - The unphased 5' to 3' sequence of nucleotides seen at a subset of the polymorphic sites examined herein in a locus on a pair of homologous chromosomes in a single individual.

**Genotyping** - A process for determining a genotype of an individual.

**Haplotype** - A 5' to 3' sequence of nucleotides found at one or more polymorphic sites in a locus on a single chromosome from a single individual. As used herein, haplotype includes a full-haplotype and/or a sub-haplotype as described below.

**Full-haplotype** - The 5' to 3' sequence of nucleotides found at all polymorphic sites examined herein in a locus on a single chromosome from a single individual.

**Sub-haplotype** - The 5' to 3' sequence of nucleotides seen at a subset of the polymorphic sites examined herein in a locus on a single chromosome from a single individual.

**Haplotype pair** - The two haplotypes found for a locus in a single individual.

**Haplotyping** - A process for determining one or more haplotypes in an individual and includes use of family pedigrees, molecular techniques and/or statistical inference.

**Haplotype data** - Information concerning one or more of the following for a specific gene: a listing of the haplotype pairs in each individual in a population; a listing of the different haplotypes in a population; frequency of each haplotype in that or other populations, and any known associations between one or more haplotypes and a trait.

**Isoform** - A particular form of a gene, mRNA, cDNA, coding sequence or the protein encoded thereby, distinguished from other forms by its particular sequence and/or structure.

**Isogene** - One of the isoforms (e.g., alleles) of a gene found in a population. An isogene (or allele) contains all of the polymorphisms present in the particular iso form of the gene.

**Isolated** - As applied to a biological molecule such as RNA, DNA, oligonucleotide, or protein,

isolated means the molecule is substantially free of other biological molecules such as nucleic acids, proteins, lipids, carbohydrates, or other material such as cellular debris and growth media. Generally, the term "isolated" is not intended to refer to a complete absence of such material or to absence of water, buffers, or salts, unless they are present in amounts that substantially interfere with the methods of the present invention.

**Locus** - A location on a chromosome or DNA molecule corresponding to a gene or a physical or phenotypic feature, where physical features include polymorphic sites.

**Naturally-occurring** - A term used to designate that the object it is applied to, e.g., naturally-occurring polynucleotide or polypeptide, can be isolated from a source in nature and which has not been intentionally modified by man.

**Nucleotide pair** - The nucleotides found at a polymorphic site on the two copies of a chromosome from an individual.

**Phased** - As applied to a sequence of nucleotide pairs for two or more polymorphic sites in a locus, phased means the combination of nucleotides present at those polymorphic sites on a single copy of the locus is known.

**Polymorphic site (PS)** - A position on a chromosome or DNA molecule at which at least two alternative sequences are found in a population.

**Polymorphic variant (variant)** - A gene, mRNA, cDNA, polypeptide, protein or peptide whose nucleotide or amino acid sequence varies from a reference sequence due to the presence of a polymorphism in the gene.

**Polymorphism** - The sequence variation observed in an individual at a polymorphic site. Polymorphisms include nucleotide substitutions, insertions, deletions and microsatellites and may, but need not, result in detectable differences in gene expression or protein function.

**Polymorphism data** - Information concerning one or more of the following for a specific gene: location of polymorphic sites; sequence variation at those sites; frequency of polymorphisms in one or more populations; the different genotypes and/or haplotypes determined for the gene; frequency of one or more of these genotypes and/or haplotypes in one or more populations; any known association(s) between a trait and a genotype or a haplotype for the gene.

**Polymorphism Database** - A collection of polymorphism data arranged in a systematic or methodical way and capable of being individually accessed by electronic or other means.

**Polynucleotide** - A nucleic acid molecule comprised of single-stranded RNA or DNA or comprised of complementary, double-stranded DNA.

**Population Group** - A group of individuals sharing a common ethnogeographic origin.

**Reference Population** - A group of subjects or individuals who are predicted to be representative

of the genetic variation found in the general population. Typically, the reference population represents the genetic variation in the population at a certainty level of at least 85%, preferably at least 90%, more preferably at least 95% and even more preferably at least 99%.

5       **Single Nucleotide Polymorphism (SNP)** – Typically, the specific pair of nucleotides observed at a single polymorphic site. In rare cases, three or four nucleotides may be found.

**Subject** – A human individual whose genotypes or haplotypes or response to treatment or disease state are to be determined.

**Treatment** – A stimulus administered internally or externally to a subject.

10       **Unphased** – As applied to a sequence of nucleotide pairs for two or more polymorphic sites in a locus, unphased means the combination of nucleotides present at those polymorphic sites on a single copy of the locus is not known.

      As discussed above, information on the identity of genotypes and haplotypes for the TNFRSF11B gene of any particular individual as well as the frequency of such genotypes and haplotypes in any particular population of individuals is useful for a variety of drug discovery and development applications.  
15       Thus, the invention also provides compositions and methods for detecting the novel TNFRSF11B polymorphisms, haplotypes and haplotype pairs identified herein.

      The compositions comprise at least one oligonucleotide for detecting the variant nucleotide or nucleotide pair located at a novel TNFRSF11B polymorphic site in one copy or two copies of the TNFRSF11B gene. Such oligonucleotides are referred to herein as TNFRSF11B haplotyping  
20       oligonucleotides or genotyping oligonucleotides, respectively, and collectively as TNFRSF11B oligonucleotides. In one embodiment, a TNFRSF11B haplotyping or genotyping oligonucleotide is a probe or primer capable of hybridizing to a target region that contains, or that is located close to, one of the novel polymorphic sites described herein.

      As used herein, the term “oligonucleotide” refers to a polynucleotide molecule having less than  
25       about 100 nucleotides. A preferred oligonucleotide of the invention is 10 to 35 nucleotides long. More preferably, the oligonucleotide is between 15 and 30, and most preferably, between 20 and 25 nucleotides in length. The exact length of the oligonucleotide will depend on many factors that are routinely considered and practiced by the skilled artisan. The oligonucleotide may be comprised of any phosphorylation state of ribonucleotides, deoxyribonucleotides, and acyclic nucleotide derivatives, and  
30       other functionally equivalent derivatives. Alternatively, oligonucleotides may have a phosphate-free backbone, which may be comprised of linkages such as carboxymethyl, acetamidate, carbamate, polyamide (peptide nucleic acid (PNA)) and the like (Varma, R. in Molecular Biology and Biotechnology, A Comprehensive Desk Reference, Ed. R. Meyers, VCH Publishers, Inc. (1995), pages 617-620).  
      Oligonucleotides of the invention may be prepared by chemical synthesis using any suitable methodology

known in the art, or may be derived from a biological sample, for example, by restriction digestion. The oligonucleotides may be labeled, according to any technique known in the art, including use of radiolabels, fluorescent labels, enzymatic labels, proteins, haptens, antibodies, sequence tags and the like.

Haplotyping or genotyping oligonucleotides of the invention must be capable of specifically hybridizing to a target region of a TNFRSF11B polynucleotide. Preferably, the target region is located in a TNFRSF11B isogene. As used herein, specific hybridization means the oligonucleotide forms an anti-parallel double-stranded structure with the target region under certain hybridizing conditions, while failing to form such a structure when incubated with another region in the TNFRSF11B polynucleotide or with a non-TNFRSF11B polynucleotide under the same hybridizing conditions. Preferably, the oligonucleotide specifically hybridizes to the target region under conventional high stringency conditions. The skilled artisan can readily design and test oligonucleotide probes and primers suitable for detecting polymorphisms in the TNFRSF11B gene using the polymorphism information provided herein in conjunction with the known sequence information for the TNFRSF11B gene and routine techniques.

A nucleic acid molecule such as an oligonucleotide or polynucleotide is said to be a "perfect" or "complete" complement of another nucleic acid molecule if every nucleotide of one of the molecules is complementary to the nucleotide at the corresponding position of the other molecule. A nucleic acid molecule is "substantially complementary" to another molecule if it hybridizes to that molecule with sufficient stability to remain in a duplex form under conventional low-stringency conditions. Conventional hybridization conditions are described, for example, by Sambrook J. et al., in *Molecular Cloning, A Laboratory Manual*, 2<sup>nd</sup> Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989) and by Haymes, B.D. et al. in *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985). While perfectly complementary oligonucleotides are preferred for detecting polymorphisms, departures from complete complementarity are contemplated where such departures do not prevent the molecule from specifically hybridizing to the target region. For example, an oligonucleotide primer may have a non-complementary fragment at its 5' end, with the remainder of the primer being complementary to the target region. Alternatively, non-complementary nucleotides may be interspersed into the probe or primer as long as the resulting probe or primer is still capable of specifically hybridizing to the target region.

Preferred haplotyping or genotyping oligonucleotides of the invention are allele-specific oligonucleotides. As used herein, the term allele-specific oligonucleotide (ASO) means an oligonucleotide that is able, under sufficiently stringent conditions, to hybridize specifically to one allele of a gene, or other locus, at a target region containing a polymorphic site while not hybridizing to the corresponding region in another allele(s). As understood by the skilled artisan, allele-specificity will depend upon a variety of readily optimized stringency conditions, including salt and formamide concentrations, as well as

temperatures for both the hybridization and washing steps. Examples of hybridization and washing conditions typically used for ASO probes are found in Kogan et al., "Genetic Prediction of Hemophilia A" in PCR Protocols, A Guide to Methods and Applications, Academic Press, 1990 and Ruaño et al., 87 *Proc. Natl. Acad. Sci. USA* 6296-6300, 1990. Typically, an ASO will be perfectly complementary to one allele while containing a single mismatch for another allele.

Allele-specific oligonucleotides of the invention include ASO probes and ASO primers. ASO probes which usually provide good discrimination between different alleles are those in which a central position of the oligonucleotide probe aligns with the polymorphic site in the target region (e.g., approximately the 7<sup>th</sup> or 8<sup>th</sup> position in a 15mer, the 8<sup>th</sup> or 9<sup>th</sup> position in a 16mer, and the 10<sup>th</sup> or 11<sup>th</sup> position in a 20mer). An ASO primer of the invention has a 3' terminal nucleotide, or preferably a 3' penultimate nucleotide, that is complementary to only one nucleotide of a particular SNP, thereby acting as a primer for polymerase-mediated extension only if the allele containing that nucleotide is present. ASO probes and primers hybridizing to either the coding or noncoding strand are contemplated by the invention. ASO probes and primers listed below use the appropriate nucleotide symbol (R= G or A, Y= T or C, M= A or C, K= G or T, S= G or C, and W= A or T; WIPO standard ST.25) at the position of the polymorphic site to represent that the ASO contains either of the two alternative allelic variants observed at that polymorphic site.

A preferred ASO probe for detecting TNFRSF11B gene polymorphisms comprises a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

- 20 GATCTTGKCTGGATC (SEQ ID NO:4) and its complement,  
CCACCGCYCCACCCC (SEQ ID NO:5) and its complement,  
TCCCTGGKGGATCCT (SEQ ID NO:6) and its complement,  
GCGTTAAYCCTGGAG (SEQ ID NO:7) and its complement,  
CCTGGGCYAGCCGAC (SEQ ID NO:8) and its complement,  
25 GGGAGAARGCTCCAC (SEQ ID NO:9) and its complement,  
CCTTTTAMGCTGCAA (SEQ ID NO:10) and its complement,  
GCTGGTAYGTGTCAA (SEQ ID NO:11) and its complement,  
AGGACCAYTGCTCAG (SEQ ID NO:12) and its complement,  
AACATAAYAGTAGCA (SEQ ID NO:13) and its complement,  
30 TATTTTCYGTAGGAA (SEQ ID NO:14) and its complement,  
CATTTTARCATATTT (SEQ ID NO:15) and its complement,  
AAGTAAAYGCAGAGA (SEQ ID NO:16) and its complement,  
AGAGGATRAAACGGC (SEQ ID NO:17) and its complement,  
TGAAGTTRTGGAAC (SEQ ID NO:18) and its complement,  
35 GTATGATMATCTAAA (SEQ ID NO:19) and its complement,  
AAACAGCRTGCAGCG (SEQ ID NO:20) and its complement, and  
TCAGAAGYTATTTTT (SEQ ID NO:21); and its complement.

- 40 A preferred ASO primer for detecting TNFRSF11B gene polymorphisms comprises a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

5 CGTCCGGATCTTGKC (SEQ ID NO:22); GAGTCCGATCCAGMC (SEQ ID NO:23);  
 CAGACACCACCGCYC (SEQ ID NO:24); GCGTGAGGGGTGGRG (SEQ ID NO:25);  
 CCCACCTCCCTGGKG (SEQ ID NO:26); GCGGAAAGGATCCMC (SEQ ID NO:27);  
 CTGAAAGCGTTAAYC (SEQ ID NO:28); AGAAAGCTCCAGGRT (SEQ ID NO:29);  
 TAAGTCCCTGGGCTA (SEQ ID NO:30); GCACCCGTCGGCTRG (SEQ ID NO:31);  
 CCGGCGGGGAGAARG (SEQ ID NO:32); GAGCGAGTGGAGCYT (SEQ ID NO:33);  
 GGGTGTCTTTTAMG (SEQ ID NO:34); GGAACCTTGCAGCKT (SEQ ID NO:35);  
 GTGCAAGCTGGTAYG (SEQ ID NO:36); TGCACATTGACACRT (SEQ ID NO:37);  
 10 TTCAAAGGACCAYT (SEQ ID NO:38); ATTCCTCTGAGCART (SEQ ID NO:39);  
 TTGTGCAACATAAYA (SEQ ID NO:40); TTTTACTGCTACTRT (SEQ ID NO:41);  
 GCTTGGTATTTTCYG (SEQ ID NO:42); CTGGGGTTCCTACRG (SEQ ID NO:43);  
 ACTTTGCATTTTARC (SEQ ID NO:44); AAGATAAAATATGYT (SEQ ID NO:45);  
 GCACCAAAGTAAAYG (SEQ ID NO:46); CTACACTCTCTGCRT (SEQ ID NO:47);  
 15 GTGTAGAGAGGATRA (SEQ ID NO:48); TGTGTTGCCGTTTYA (SEQ ID NO:49);  
 AGCTGCTGAAGTTRT (SEQ ID NO:50); TTTGATGTTTCCAYA (SEQ ID NO:51);  
 TCCAAGGTATGATMA (SEQ ID NO:52); TTTTATTTTAGATKA (SEQ ID NO:53);  
 CTGTGAAAACAGCRT (SEQ ID NO:54); ATGTGCCGCTGCAYG (SEQ ID NO:55);  
 20 ATTGATCAGAAGYT (SEQ ID NO:56); and ATTTCTAAAAATARC (SEQ ID NO:57).

Other oligonucleotides of the invention hybridize to a target region located one to several nucleotides downstream of one of the novel polymorphic sites identified herein. Such oligonucleotides are useful in polymerase-mediated primer extension methods for detecting one of the novel polymorphisms described herein and therefore such oligonucleotides are referred to herein as "primer-extension oligonucleotides". In a preferred embodiment, the 3'-terminus of a primer-extension oligonucleotide is a deoxynucleotide complementary to the nucleotide located immediately adjacent to the polymorphic site.

A particularly preferred oligonucleotide primer for detecting TNFRSF11B gene polymorphisms by primer extension terminates in a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

30 CCGGATCTTG (SEQ ID NO:58); TCCGATCCAG (SEQ ID NO:59);  
 ACACCACCGC (SEQ ID NO:60); TGAGGGGTGG (SEQ ID NO:61);  
 ACCTCCCTGG (SEQ ID NO:62); GAAAGGATCC (SEQ ID NO:63);  
 AAAGCGTTAA (SEQ ID NO:64); AAGCTCCAGG (SEQ ID NO:65);  
 GTCCCTGGGC (SEQ ID NO:66); CCCGTCGGCT (SEQ ID NO:67);  
 35 GCGGGGAGAA (SEQ ID NO:68); CGAGTGGAGC (SEQ ID NO:69);  
 TGTCCTTTTA (SEQ ID NO:70); ACTTTGCAGC (SEQ ID NO:71);  
 CAAGCTGGTA (SEQ ID NO:72); ACATTGACAC (SEQ ID NO:73);  
 CAAAGGACCA (SEQ ID NO:74); CCTCTGAGCA (SEQ ID NO:75);  
 TGCAACATAA (SEQ ID NO:76); TACTGCTACT (SEQ ID NO:77);  
 TGGTATTTTC (SEQ ID NO:78); GGGTTCCTAC (SEQ ID NO:79);  
 40 TTGCATTTTA (SEQ ID NO:80); ATAAAATATG (SEQ ID NO:81);  
 CCAAAGTAAA (SEQ ID NO:82); CACTCTCTGC (SEQ ID NO:83);  
 TAGAGAGGAT (SEQ ID NO:84); GTTGCCGTTT (SEQ ID NO:85);  
 TGCTGAAGTT (SEQ ID NO:86); GATGTTTCCA (SEQ ID NO:87);  
 AAGGTATGAT (SEQ ID NO:88); TATTTTAGAT (SEQ ID NO:89);  
 45 TGAAAACAGC (SEQ ID NO:90); TGCCGCTGCA (SEQ ID NO:91);  
 GTATCAGAAG (SEQ ID NO:92); and TCTAAAAATA (SEQ ID NO:93).

In some embodiments, a composition contains two or more differently labeled TNFRSF11B oligonucleotides for simultaneously probing the identity of nucleotides or nucleotide pairs at two or more polymorphic sites. It is also contemplated that primer compositions may contain two or more sets of allele-specific primer pairs to allow simultaneous targeting and amplification of two or more regions containing a polymorphic site.

TNFRSF11B oligonucleotides of the invention may also be immobilized on or synthesized on a solid surface such as a microchip, bead, or glass slide (see, e.g., WO 98/20020 and WO 98/20019). Such immobilized oligonucleotides may be used in a variety of polymorphism detection assays, including but not limited to probe hybridization and polymerase extension assays. Immobilized TNFRSF11B oligonucleotides of the invention may comprise an ordered array of oligonucleotides designed to rapidly screen a DNA sample for polymorphisms in multiple genes at the same time.

In another embodiment, the invention provides a kit comprising at least two TNFRSF11B oligonucleotides packaged in separate containers. The kit may also contain other components such as hybridization buffer (where the oligonucleotides are to be used as a probe) packaged in a separate container. Alternatively, where the oligonucleotides are to be used to amplify a target region, the kit may contain, packaged in separate containers, a polymerase and a reaction buffer optimized for primer extension mediated by the polymerase, such as PCR.

The above described oligonucleotide compositions and kits are useful in methods for genotyping and/or haplotyping the TNFRSF11B gene in an individual. As used herein, the terms "TNFRSF11B genotype" and "TNFRSF11B haplotype" mean the genotype or haplotype contains the nucleotide pair or nucleotide, respectively, that is present at one or more of the novel polymorphic sites described herein and may optionally also include the nucleotide pair or nucleotide present at one or more additional polymorphic sites in the TNFRSF11B gene. The additional polymorphic sites may be currently known polymorphic sites or sites that are subsequently discovered.

One embodiment of a genotyping method of the invention involves examining both copies of the individual's TNFRSF11B gene, or a fragment thereof, to identify the nucleotide pair at one or more polymorphic sites selected from the group consisting of PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19 in the two copies to assign a TNFRSF11B genotype to the individual. In some embodiments, "examining a gene" may include examining one or more of: DNA containing the gene, mRNA transcripts thereof, or cDNA copies thereof. As will be readily understood by the skilled artisan, the two "copies" of a gene, mRNA or cDNA (or fragment of such TNFRSF11B molecules) in an individual may be the same allele or may be different alleles. In a preferred embodiment of the method for assigning a TNFRSF11B genotype, the identity of the nucleotide pair at PS5 is also determined. In another embodiment, a genotyping method of the invention comprises

determining the identity of the nucleotide pair at each of PS1-PS19.

One method of examining both copies of the individual's TNFRSF11B gene is by isolating from the individual a nucleic acid sample comprising the two copies of the TNFRSF11B gene, mRNA transcripts thereof or cDNA copies thereof, or a fragment of any of the foregoing, that are present in the individual. Typically, the nucleic acid sample is isolated from a biological sample taken from the individual, such as a blood sample or tissue sample. Suitable tissue samples include whole blood, semen, saliva, tears, urine, fecal material, sweat, buccal, skin and hair. The nucleic acid sample may be comprised of genomic DNA, mRNA, or cDNA and, in the latter two cases, the biological sample must be obtained from a tissue in which the TNFRSF11B gene is expressed. Furthermore it will be understood by the skilled artisan that mRNA or cDNA preparations would not be used to detect polymorphisms located in introns or in 5' and 3' untranslated regions if not present in the mRNA or cDNA. If a TNFRSF11B gene fragment is isolated, it must contain the polymorphic site(s) to be genotyped.

One embodiment of a haplotyping method of the invention comprises examining one copy of the individual's TNFRSF11B gene, or a fragment thereof, to identify the nucleotide at one or more polymorphic sites selected from the group consisting of PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19 in that copy to assign a TNFRSF11B haplotype to the individual. In some embodiments, "examining a gene" may include examining one or more of: DNA containing the gene, mRNA transcripts thereof, or cDNA copies thereof. One method of examining one copy of the individual's TNFRSF11B gene is by isolating from the individual a nucleic acid sample containing only one of the two copies of the TNFRSF11B gene, mRNA or cDNA, or a fragment of such TNFRSF11B molecules, that is present in the individual and determining in that copy the identity of the nucleotide at one or more polymorphic sites selected from the group consisting of PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19.

The nucleic acid used in the above haplotyping methods of the invention may be isolated using any method capable of separating the two copies of the TNFRSF11B gene or fragment such as one of the methods described above for preparing TNFRSF11B isogenes, with targeted *in vivo* cloning being the preferred approach. As will be readily appreciated by those skilled in the art, any individual clone will typically only provide haplotype information on one of the two TNFRSF11B gene copies present in an individual. If haplotype information is desired for the individual's other copy, additional TNFRSF11B clones will usually need to be examined. Typically, at least five clones should be examined to have more than a 90% probability of haplotyping both copies of the TNFRSF11B gene in an individual. In some cases, however, once the haplotype for one TNFRSF11B allele is directly determined, the haplotype for the other allele may be inferred if the individual has a known genotype for the polymorphic sites of interest or if the haplotype frequency or haplotype pair frequency for the individual's population group is known. In

some embodiments, the TNFRSF11B haplotype is assigned to the individual by also identifying the nucleotide at PS5. In a particularly preferred embodiment, the nucleotide at each of PS1-PS19 is identified.

In another embodiment, the haplotyping method comprises determining whether an individual has one or more of the TNFRSF11B haplotypes shown in Table 5. This can be accomplished by identifying, for one or both copies of the individual's TNFRSF11B gene, the phased sequence of nucleotides present at each of PS1-PS19. This identifying step does not necessarily require that each of PS1-PS19 be directly examined. Typically only a subset of PS1-PS19 will need to be directly examined to assign to an individual one or more of the haplotypes shown in Table 5. This is because at least one polymorphic site in a gene is frequently in strong linkage disequilibrium with one or more other polymorphic sites in that gene (Drysdales, CM et al. 2000 *PNAS* 97:10483-10488; Rieder MJ et al. 1999 *Nature Genetics* 22:59-62). Two sites are said to be in linkage disequilibrium if the presence of a particular variant at one site enhances the predictability of another variant at the second site (Stephens, JC 1999, *Mol. Diag.* 4:309-317). Techniques for determining whether any two polymorphic sites are in linkage disequilibrium are well-known in the art (Weir B.S. 1996 *Genetic Data Analysis II*, Sinauer Associates, Inc. Publishers, Sunderland, MA). In addition, Johnson et al. (2001 *Nature Genetics* 29: 233-237) presented one possible method for selection of subsets of polymorphic sites suitable for identifying known haplotypes.

In another embodiment of a haplotyping method of the invention, a TNFRSF11B haplotype pair is determined for an individual by identifying the phased sequence of nucleotides at one or more polymorphic sites selected from the group consisting of PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19 in each copy of the TNFRSF11B gene that is present in the individual. In a particularly preferred embodiment, the haplotyping method comprises identifying the phased sequence of nucleotides at each of PS1-PS19 in each copy of the TNFRSF11B gene.

When haplotyping both copies of the gene, the identifying step is preferably performed with each copy of the gene being placed in separate containers. However, it is also envisioned that if the two copies are labeled with different tags, or are otherwise separately distinguishable or identifiable, it could be possible in some cases to perform the method in the same container. For example, if first and second copies of the gene are labeled with different first and second fluorescent dyes, respectively, and an allele-specific oligonucleotide labeled with yet a third different fluorescent dye is used to assay the polymorphic site(s), then detecting a combination of the first and third dyes would identify the polymorphism in the first gene copy while detecting a combination of the second and third dyes would identify the polymorphism in the second gene copy.

In both the genotyping and haplotyping methods, the identity of a nucleotide (or nucleotide pair) at a polymorphic site(s) may be determined by amplifying a target region(s) containing the polymorphic

site(s) directly from one or both copies of the TNFRSF11B gene, or a fragment thereof, and the sequence of the amplified region(s) determined by conventional methods. It will be readily appreciated by the skilled artisan that only one nucleotide will be detected at a polymorphic site in individuals who are homozygous at that site, while two different nucleotides will be detected if the individual is heterozygous for that site. The polymorphism may be identified directly, known as positive-type identification, or by inference, referred to as negative-type identification. For example, where a SNP is known to be guanine and cytosine in a reference population, a site may be positively determined to be either guanine or cytosine for an individual homozygous at that site, or both guanine and cytosine, if the individual is heterozygous at that site. Alternatively, the site may be negatively determined to be not guanine (and thus cytosine/cytosine) or not cytosine (and thus guanine/guanine).

The target region(s) may be amplified using any oligonucleotide-directed amplification method, including but not limited to polymerase chain reaction (PCR) (U.S. Patent No. 4,965,188), ligase chain reaction (LCR) (Barany et al., *Proc. Natl. Acad. Sci. USA* 88:189-193, 1991; WO90/01069), and oligonucleotide ligation assay (OLA) (Landegren et al., *Science* 241:1077-1080, 1988). Other known nucleic acid amplification procedures may be used to amplify the target region including transcription-based amplification systems (U.S. Patent No. 5,130,238; EP 329,822; U.S. Patent No. 5,169,766, WO89/06700) and isothermal methods (Walker et al., *Proc. Natl. Acad. Sci. USA* 89:392-396, 1992).

A polymorphism in the target region may also be assayed before or after amplification using one of several hybridization-based methods known in the art. Typically, allele-specific oligonucleotides are utilized in performing such methods. The allele-specific oligonucleotides may be used as differently labeled probe pairs, with one member of the pair showing a perfect match to one variant of a target sequence and the other member showing a perfect match to a different variant. In some embodiments, more than one polymorphic site may be detected at once using a set of allele-specific oligonucleotides or oligonucleotide pairs. Preferably, the members of the set have melting temperatures within 5°C, and more preferably within 2°C, of each other when hybridizing to each of the polymorphic sites being detected.

Hybridization of an allele-specific oligonucleotide to a target polynucleotide may be performed with both entities in solution, or such hybridization may be performed when either the oligonucleotide or the target polynucleotide is covalently or noncovalently affixed to a solid support. Attachment may be mediated, for example, by antibody-antigen interactions, poly-L-Lys, streptavidin or avidin-biotin, salt bridges, hydrophobic interactions, chemical linkages, UV cross-linking baking, etc. Allele-specific oligonucleotides may be synthesized directly on the solid support or attached to the solid support subsequent to synthesis. Solid-supports suitable for use in detection methods of the invention include substrates made of silicon, glass, plastic, paper and the like, which may be formed, for example, into wells (as in 96-well plates), slides, sheets, membranes, fibers, chips, dishes, and beads. The solid support may

be treated, coated or derivatized to facilitate the immobilization of the allele-specific oligonucleotide or target nucleic acid.

The genotype or haplotype for the TNFRSF11B gene of an individual may also be determined by hybridization of a nucleic acid sample containing one or both copies of the gene, mRNA, cDNA or fragment(s) thereof, to nucleic acid arrays and subarrays such as described in WO 95/11995. The arrays would contain a battery of allele-specific oligonucleotides representing each of the polymorphic sites to be included in the genotype or haplotype.

The identity of polymorphisms may also be determined using a mismatch detection technique, including but not limited to the RNase protection method using riboprobes (Winter et al., *Proc. Natl. Acad. Sci. USA* 82:7575, 1985; Meyers et al., *Science* 230:1242, 1985) and proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein (Modrich, *P. Ann. Rev. Genet.* 25:229-253, 1991).

Alternatively, variant alleles can be identified by single strand conformation polymorphism (SSCP) analysis (Orita et al., *Genomics* 5:874-879, 1989; Humphries et al., in *Molecular Diagnosis of Genetic Diseases*, R. Elles, ed., pp. 321-340, 1996) or denaturing gradient gel electrophoresis (DGGE) (Wartell et al., *Nucl. Acids Res.* 18:2699-2706, 1990; Sheffield et al., *Proc. Natl. Acad. Sci. USA* 86:232-236, 1989).

A polymerase-mediated primer extension method may also be used to identify the polymorphism(s). Several such methods have been described in the patent and scientific literature and include the "Genetic Bit Analysis" method (WO92/15712) and the ligase/polymerase mediated genetic bit analysis (U.S. Patent 5,679,524. Related methods are disclosed in WO91/02087, WO90/09455, WO95/17676, U.S. Patent Nos. 5,302,509, and 5,945,283. Extended primers containing a polymorphism may be detected by mass spectrometry as described in U.S. Patent No. 5,605,798. Another primer extension method is allele-specific PCR (Ruano et al., *Nucl. Acids Res.* 17:8392, 1989; Ruano et al., *Nucl. Acids Res.* 19, 6877-6882, 1991; WO 93/22456; Turki et al., *J. Clin. Invest.* 95:1635-1641, 1995). In addition, multiple polymorphic sites may be investigated by simultaneously amplifying multiple regions of the nucleic acid using sets of allele-specific primers as described in Wallace et al. (WO89/10414).

In addition, the identity of the allele(s) present at any of the novel polymorphic sites described herein may be indirectly determined by haplotyping or genotyping another polymorphic site that is in linkage disequilibrium with the polymorphic site that is of interest. Polymorphic sites in linkage disequilibrium with the presently disclosed polymorphic sites may be located in regions of the gene or in other genomic regions not examined herein. Detection of the allele(s) present at a polymorphic site in linkage disequilibrium with the novel polymorphic sites described herein may be performed by, but is not limited to, any of the above-mentioned methods for detecting the identity of the allele at a polymorphic site.

In another aspect of the invention, an individual's TNFRSF11B haplotype pair is predicted from

its TNFRSF11B genotype using information on haplotype pairs known to exist in a reference population. In its broadest embodiment, the haplotyping prediction method comprises identifying a TNFRSF11B genotype for the individual at two or more TNFRSF11B polymorphic sites described herein, accessing data containing TNFRSF11B haplotype pairs identified in a reference population, and assigning a haplotype pair to the individual that is consistent with the genotype data. In one embodiment, the reference haplotype pairs include the TNFRSF11B haplotype pairs shown in Table 4. The TNFRSF11B haplotype pair can be assigned by comparing the individual's genotype with the genotypes corresponding to the haplotype pairs known to exist in the general population or in a specific population group, and determining which haplotype pair is consistent with the genotype of the individual. In some embodiments, the comparing step may be performed by visual inspection (for example, by consulting Table 4). When the genotype of the individual is consistent with more than one haplotype pair, frequency data (such as that presented in Table 7) may be used to determine which of these haplotype pairs is most likely to be present in the individual. This determination may also be performed in some embodiments by visual inspection, for example by consulting Table 7. If a particular TNFRSF11B haplotype pair consistent with the genotype of the individual is more frequent in the reference population than others consistent with the genotype, then that haplotype pair with the highest frequency is the most likely to be present in the individual. In other embodiments, the comparison may be made by a computer-implemented algorithm with the genotype of the individual and the reference haplotype data stored in computer-readable formats. For example, as described in PCT/US01/12831, filed April 18, 2001, one computer-implemented algorithm to perform this comparison entails enumerating all possible haplotype pairs which are consistent with the genotype, accessing data containing TNFRSF11B haplotype pairs frequency data determined in a reference population to determine a probability that the individual has a possible haplotype pair, and analyzing the determined probabilities to assign a haplotype pair to the individual.

Generally, the reference population should be composed of randomly-selected individuals representing the major ethnogeographic groups of the world. A preferred reference population for use in the methods of the present invention comprises an approximately equal number of individuals from Caucasian, African-descent, Asian and Hispanic-Latino population groups with the minimum number of each group being chosen based on how rare a haplotype one wants to be guaranteed to see. For example, if one wants to have a q% chance of not missing a haplotype that exists in the population at a p% frequency of occurring in the reference population, the number of individuals (n) who must be sampled is given by  $2n = \log(1-q)/\log(1-p)$  where p and q are expressed as fractions. A preferred reference population allows the detection of any haplotype whose frequency is at least 10% with about 99% certainty and comprises about 20 unrelated individuals from each of the four population groups named above. A particularly preferred reference population includes a 3-generation family representing one or more of the four

population groups to serve as controls for checking quality of haplotyping procedures.

In a preferred embodiment, the haplotype frequency data for each ethnogeographic group is examined to determine whether it is consistent with Hardy-Weinberg equilibrium. Hardy-Weinberg equilibrium (D.L. Hartl et al., Principles of Population Genomics, Sinauer Associates (Sunderland, MA),

3<sup>rd</sup> Ed., 1997) postulates that the frequency of finding the haplotype pair  $H_1 / H_2$  is equal to

$p_{H-W}(H_1 / H_2) = 2p(H_1)p(H_2)$  if  $H_1 \neq H_2$  and  $p_{H-W}(H_1 / H_2) = p(H_1)p(H_2)$  if  $H_1 = H_2$ . A

statistically significant difference between the observed and expected haplotype frequencies could be due to one or more factors including significant inbreeding in the population group, strong selective pressure on the gene, sampling bias, and/or errors in the genotyping process. If large deviations from Hardy-

Weinberg equilibrium are observed in an ethnogeographic group, the number of individuals in that group can be increased to see if the deviation is due to a sampling bias. If a larger sample size does not reduce the difference between observed and expected haplotype pair frequencies, then one may wish to consider haplotyping the individual using a direct haplotyping method such as, for example, CLASPER System™ technology (U.S. Patent No. 5,866,404), single molecule dilution, or allele-specific long-range PCR (Michalotos-Beloin et al., *Nucleic Acids Res.* 24:4841-4843, 1996).

In one embodiment of this method for predicting a TNFRSF11B haplotype pair for an individual, the assigning step involves performing the following analysis. First, each of the possible haplotype pairs is compared to the haplotype pairs in the reference population. Generally, only one of the haplotype pairs in the reference population matches a possible haplotype pair and that pair is assigned to the individual.

Occasionally, only one haplotype represented in the reference haplotype pairs is consistent with a possible haplotype pair for an individual, and in such cases the individual is assigned a haplotype pair containing this known haplotype and a new haplotype derived by subtracting the known haplotype from the possible haplotype pair. Alternatively, the haplotype pair in an individual may be predicted from the individual's genotype for that gene using reported methods (e.g., Clark et al. 1990 *Mol Bio Evol* 7:111-22; copending PCT/US01/12831 filed April 18, 2001 ) or through a commercial haplotyping service such as offered by Genaissance Pharmaceuticals, Inc. (New Haven, CT). In rare cases, either no haplotypes in the reference population are consistent with the possible haplotype pairs, or alternatively, multiple reference haplotype pairs are consistent with the possible haplotype pairs. In such cases, the individual is preferably haplotyped using a direct molecular haplotyping method such as, for example, CLASPER System™ technology (U.S. Patent No. 5,866,404), SMD, or allele-specific long-range PCR (Michalotos-Beloin et al., *supra*).

The invention also provides a method for determining the frequency of a TNFRSF11B genotype, haplotype, or haplotype pair in a population. The method comprises, for each member of the population,

determining the genotype or the haplotype pair for the novel TNFRSF11B polymorphic sites described herein, and calculating the frequency any particular genotype, haplotype, or haplotype pair is found in the population. The population may be e.g., a reference population, a family population, a same gender population, a population group, or a trait population (e.g., a group of individuals exhibiting a trait of interest such as a medical condition or response to a therapeutic treatment).

In another aspect of the invention, frequency data for TNFRSF11B genotypes, haplotypes, and/or haplotype pairs are determined in a reference population and used in a method for identifying an association between a trait and a TNFRSF11B genotype, haplotype, or haplotype pair. The trait may be any detectable phenotype, including but not limited to susceptibility to a disease or response to a treatment.

In one embodiment, the method involves obtaining data on the frequency of the genotype(s), haplotype(s), or haplotype pair(s) of interest in a reference population as well as in a population exhibiting the trait.

Frequency data for one or both of the reference and trait populations may be obtained by genotyping or haplotyping each individual in the populations using one or more of the methods described above. The haplotypes for the trait population may be determined directly or, alternatively, by a predictive genotype to haplotype approach as described above. In another embodiment, the frequency data for the reference and/or trait populations is obtained by accessing previously determined frequency data, which may be in written or electronic form. For example, the frequency data may be present in a database that is accessible by a computer. Once the frequency data is obtained, the frequencies of the genotype(s), haplotype(s), or haplotype pair(s) of interest in the reference and trait populations are compared. In a preferred

embodiment, the frequencies of all genotypes, haplotypes, and/or haplotype pairs observed in the populations are compared. If a particular TNFRSF11B genotype, haplotype, or haplotype pair is more frequent in the trait population than in the reference population at a statistically significant amount, then the trait is predicted to be associated with that TNFRSF11B genotype, haplotype or haplotype pair. Preferably, the TNFRSF11B genotype, haplotype, or haplotype pair being compared in the trait and reference populations is selected from the full-genotypes and full-haplotypes shown in Tables 4 and 5, or from sub-genotypes and sub-haplotypes derived from these genotypes and haplotypes. Sub-genotypes useful in the invention preferably do not include the sub-genotype consisting solely of PS5.

In a preferred embodiment of the method, the trait of interest is a clinical response exhibited by a patient to some therapeutic treatment, for example, response to a drug targeting TNFRSF11B or response to a therapeutic treatment for a medical condition. As used herein, "medical condition" includes but is not limited to any condition or disease manifested as one or more physical and/or psychological symptoms for which treatment is desirable, and includes previously and newly identified diseases and other disorders. As used herein the term "clinical response" means any or all of the following: a quantitative measure of the response, no response, and/or adverse response (i.e., side effects).

10044674-010902

5 In order to deduce a correlation between clinical response to a treatment and a TNFRSF11B genotype, haplotype, or haplotype pair, it is necessary to obtain data on the clinical responses exhibited by a population of individuals who received the treatment, hereinafter the "clinical population". This clinical data may be obtained by analyzing the results of a clinical trial that has already been run and/or the clinical data may be obtained by designing and carrying out one or more new clinical trials. As used herein, the term "clinical trial" means any research study designed to collect clinical data on responses to a particular treatment, and includes but is not limited to phase I, phase II and phase III clinical trials. Standard methods are used to define the patient population and to enroll subjects.

10 It is preferred that the individuals included in the clinical population have been graded for the existence of the medical condition of interest. This is important in cases where the symptom(s) being presented by the patients can be caused by more than one underlying condition, and where treatment of the underlying conditions are not the same. An example of this would be where patients experience breathing difficulties that are due to either asthma or respiratory infections. If both sets were treated with an asthma medication, there would be a spurious group of apparent non-responders that did not actually have asthma.

15 These people would affect the ability to detect any correlation between haplotype and treatment outcome. This grading of potential patients could employ a standard physical exam or one or more lab tests. Alternatively, grading of patients could use haplotyping for situations where there is a strong correlation between haplotype pair and disease susceptibility or severity.

20 The therapeutic treatment of interest is administered to each individual in the trial population and each individual's response to the treatment is measured using one or more predetermined criteria. It is contemplated that in many cases, the trial population will exhibit a range of responses and that the investigator will choose the number of responder groups (e.g., low, medium, high) made up by the various responses. In addition, the TNFRSF11B gene for each individual in the trial population is genotyped and/or haplotyped, which may be done before or after administering the treatment.

25 After both the clinical and polymorphism data have been obtained, correlations between individual response and TNFRSF11B genotype or haplotype content are created. Correlations may be produced in several ways. In one method, individuals are grouped by their TNFRSF11B genotype or haplotype (or haplotype pair) (also referred to as a polymorphism group), and then the averages and standard deviations of clinical responses exhibited by the members of each polymorphism group are calculated.

30 These results are then analyzed to determine if any observed variation in clinical response between polymorphism groups is statistically significant. Statistical analysis methods which may be used are described in L.D. Fisher and G. vanBelle, "Biostatistics: A Methodology for the Health Sciences", Wiley-Interscience (New York) 1993. This analysis may also include a regression calculation of which polymorphic sites in the TNFRSF11B gene give the most significant contribution to the differences in

phenotype. One regression model useful in the invention is described in WO 01/01218, entitled "Methods for Obtaining and Using Haplotype Data".

A second method for finding correlations between TNFRSF11B haplotype content and clinical responses uses predictive models based on error-minimizing optimization algorithms. One of many possible optimization algorithms is a genetic algorithm (R. Judson, "Genetic Algorithms and Their Uses in Chemistry" in Reviews in Computational Chemistry, Vol. 10, pp. 1-73, K. B. Lipkowitz and D. B. Boyd, eds. (VCH Publishers, New York, 1997). Simulated annealing (Press et al., "Numerical Recipes in C: The Art of Scientific Computing", Cambridge University Press (Cambridge) 1992, Ch. 10), neural networks (E. Rich and K. Knight, "Artificial Intelligence", 2<sup>nd</sup> Edition (McGraw-Hill, New York, 1991, Ch. 18), standard gradient descent methods (Press et al., *supra*, Ch. 10), or other global or local optimization approaches (see discussion in Judson, *supra*) could also be used. Preferably, the correlation is found using a genetic algorithm approach as described in WO 01/01218.

Correlations may also be analyzed using analysis of variation (ANOVA) techniques to determine how much of the variation in the clinical data is explained by different subsets of the polymorphic sites in the TNFRSF11B gene. As described in WO 01/01218, ANOVA is used to test hypotheses about whether a response variable is caused by or correlated with one or more traits or variables that can be measured (Fisher and vanBelle, *supra*, Ch. 10).

From the analyses described above, a mathematical model may be readily constructed by the skilled artisan that predicts clinical response as a function of TNFRSF11B genotype or haplotype content. Preferably, the model is validated in one or more follow-up clinical trials designed to test the model.

The identification of an association between a clinical response and a genotype or haplotype (or haplotype pair) for the TNFRSF11B gene may be the basis for designing a diagnostic method to determine those individuals who will or will not respond to the treatment, or alternatively, will respond at a lower level and thus may require more treatment, i.e., a greater dose of a drug. The diagnostic method may take one of several forms: for example, a direct DNA test (i.e., genotyping or haplotyping one or more of the polymorphic sites in the TNFRSF11B gene), a serological test, or a physical exam measurement. The only requirement is that there be a good correlation between the diagnostic test results and the underlying TNFRSF11B genotype or haplotype that is in turn correlated with the clinical response. In a preferred embodiment, this diagnostic method uses the predictive haplotyping method described above.

In another embodiment, the invention provides an isolated polynucleotide comprising a polymorphic variant of the TNFRSF11B gene or a fragment of the gene which contains at least one of the novel polymorphic sites described herein. The nucleotide sequence of a variant TNFRSF11B gene is identical to the reference genomic sequence for those portions of the gene examined, as described in the Examples below, except that it comprises a different nucleotide at one or more of the novel polymorphic

sites PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18 and PS19, and may also comprise an additional polymorphism of cytosine at PS5. Similarly, the nucleotide sequence of a variant fragment of the TNFRSF11B gene is identical to the corresponding portion of the reference sequence except for having a different nucleotide at one or more of the novel polymorphic sites described herein. Thus, the invention specifically does not include polynucleotides comprising a nucleotide sequence identical to the reference sequence of the TNFRSF11B gene, which is defined by haplotype 19, (or other reported TNFRSF11B sequences) or to portions of the reference sequence (or other reported TNFRSF11B sequences), except for the haplotyping and genotyping oligonucleotides described above.

The location of a polymorphism in a variant TNFRSF11B gene or fragment is preferably identified by aligning its sequence against SEQ ID NO:1. The polymorphism is selected from the group consisting of thymine at PS1, thymine at PS2, thymine at PS3, cytosine at PS4, thymine at PS6, adenine at PS7, adenine at PS8, thymine at PS9, cytosine at PS10, cytosine at PS11, thymine at PS12, adenine at PS13, thymine at PS14, guanine at PS15, guanine at PS16, cytosine at PS17, adenine at PS18 and cytosine at PS19. In a preferred embodiment, the polymorphic variant comprises a naturally-occurring isogene of the TNFRSF11B gene which is defined by any one of haplotypes 1- 18 and 20 - 22 shown in Table 5 below.

Polymorphic variants of the invention may be prepared by isolating a clone containing the TNFRSF11B gene from a human genomic library. The clone may be sequenced to determine the identity of the nucleotides at the novel polymorphic sites described herein. Any particular variant or fragment thereof, that is claimed herein could be prepared from this clone by performing *in vitro* mutagenesis using procedures well-known in the art. Any particular TNFRSF11B variant or fragment thereof may also be prepared using synthetic or semi-synthetic methods known in the art.

TNFRSF11B isogenes, or fragments thereof, may be isolated using any method that allows separation of the two "copies" of the TNFRSF11B gene present in an individual, which, as readily understood by the skilled artisan, may be the same allele or different alleles. Separation methods include targeted *in vivo* cloning (TIVC) in yeast as described in WO 98/01573, U.S. Patent No. 5,866,404, and U.S. Patent No. 5,972,614. Another method, which is described in U.S. Patent No. 5,972,614, uses an allele specific oligonucleotide in combination with primer extension and exonuclease degradation to generate hemizygous DNA targets. Yet other methods are single molecule dilution (SMD) as described in Ruaño et al., *Proc. Natl. Acad. Sci.* 87:6296-6300, 1990; and allele specific PCR (Ruaño et al., 1989, *supra*; Ruaño et al., 1991, *supra*; Michalatos-Beloin et al., *supra*).

The invention also provides TNFRSF11B genome anthologies, which are collections of at least two TNFRSF11B isogenes found in a given population. The population may be any group of at least two individuals, including but not limited to a reference population, a population group, a family population, a

clinical population, and a same gender population. A TNFRSF11B genome anthology may comprise individual TNFRSF11B isogenes stored in separate containers such as microtest tubes, separate wells of a microtitre plate and the like. Alternatively, two or more groups of the TNFRSF11B isogenes in the anthology may be stored in separate containers. Individual isogenes or groups of such isogenes in a genome anthology may be stored in any convenient and stable form, including but not limited to in buffered solutions, as DNA precipitates, freeze-dried preparations and the like. A preferred TNFRSF11B genome anthology of the invention comprises a set of isogenes defined by the haplotypes shown in Table 5 below.

An isolated polynucleotide containing a polymorphic variant nucleotide sequence of the invention may be operably linked to one or more expression regulatory elements in a recombinant expression vector capable of being propagated and expressing the encoded TNFRSF11B protein in a prokaryotic or a eukaryotic host cell. Examples of expression regulatory elements which may be used include, but are not limited to, the lac system, operator and promoter regions of phage lambda, yeast promoters, and promoters derived from vaccinia virus, adenovirus, retroviruses, or SV40. Other regulatory elements include, but are not limited to, appropriate leader sequences, termination codons, polyadenylation signals, and other sequences required for the appropriate transcription and subsequent translation of the nucleic acid sequence in a given host cell. Of course, the correct combinations of expression regulatory elements will depend on the host system used. In addition, it is understood that the expression vector contains any additional elements necessary for its transfer to and subsequent replication in the host cell. Examples of such elements include, but are not limited to, origins of replication and selectable markers. Such expression vectors are commercially available or are readily constructed using methods known to those in the art (e.g., F. Ausubel et al., 1987, in "Current Protocols in Molecular Biology", John Wiley and Sons, New York, New York). Host cells which may be used to express the variant TNFRSF11B sequences of the invention include, but are not limited to, eukaryotic and mammalian cells, such as animal, plant, insect and yeast cells, and prokaryotic cells, such as *E. coli*, or algal cells as known in the art. The recombinant expression vector may be introduced into the host cell using any method known to those in the art including, but not limited to, microinjection, electroporation, particle bombardment, transduction, and transfection using DEAE-dextran, lipofection, or calcium phosphate (see e.g., Sambrook et al. (1989) in "Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Press, Plainview, New York). In a preferred aspect, eukaryotic expression vectors that function in eukaryotic cells, and preferably mammalian cells, are used. Non-limiting examples of such vectors include vaccinia virus vectors, adenovirus vectors, herpes virus vectors, and baculovirus transfer vectors. Preferred eukaryotic cell lines include COS cells, CHO cells, HeLa cells, NIH/3T3 cells, and embryonic stem cells (Thomson, J. A. et al., 1998 *Science* 282:1145-1147). Particularly preferred host cells are mammalian cells.

As will be readily recognized by the skilled artisan, expression of polymorphic variants of the TNFRSF11B gene will produce TNFRSF11B mRNAs varying from each other at any polymorphic site retained in the spliced and processed mRNA molecules. These mRNAs can be used for the preparation of a TNFRSF11B cDNA comprising a nucleotide sequence which is a polymorphic variant of the TNFRSF11B reference coding sequence shown in Figure 2. Thus, the invention also provides TNFRSF11B mRNAs and corresponding cDNAs which comprise a nucleotide sequence that is identical to SEQ ID NO:2 (Fig. 2) (or its corresponding RNA sequence) for those regions of SEQ ID NO:2 that correspond to the examined portions of the TNFRSF11B gene (as described in the Examples below), except for having one or more polymorphisms selected from the group consisting of thymine at a position corresponding to nucleotide 699, guanine at a position corresponding to nucleotide 720, guanine at a position corresponding to nucleotide 768, adenine at a position corresponding to nucleotide 841 and cytosine at a position corresponding to nucleotide 1150, and may also comprise an additional polymorphism of cytosine at a position corresponding to nucleotide 9. A particularly preferred polymorphic cDNA variant comprises the coding sequence of a TNFRSF11B isogene defined by any one of haplotypes 1-3, 7-14, 18, 21 and 22. Fragments of these variant mRNAs and cDNAs are included in the scope of the invention, provided they contain one or more of the novel polymorphisms described herein. The invention specifically excludes polynucleotides identical to previously identified TNFRSF11B mRNAs or cDNAs, and previously described fragments thereof. Polynucleotides comprising a variant TNFRSF11B RNA or DNA sequence may be isolated from a biological sample using well-known molecular biological procedures or may be chemically synthesized.

As used herein, a polymorphic variant of a TNFRSF11B gene, mRNA or cDNA fragment comprises at least one novel polymorphism identified herein and has a length of at least 10 nucleotides and may range up to the full length of the gene. Preferably, such fragments are between 100 and 3000 nucleotides in length, and more preferably between 200 and 2000 nucleotides in length, and most preferably between 500 and 1000 nucleotides in length.

In describing the TNFRSF11B polymorphic sites identified herein, reference is made to the sense strand of the gene for convenience. However, as recognized by the skilled artisan, nucleic acid molecules containing the TNFRSF11B gene or cDNA may be complementary double stranded molecules and thus reference to a particular site on the sense strand refers as well to the corresponding site on the complementary antisense strand. Thus, reference may be made to the same polymorphic site on either strand and an oligonucleotide may be designed to hybridize specifically to either strand at a target region containing the polymorphic site. Thus, the invention also includes single-stranded polynucleotides which are complementary to the sense strand of the TNFRSF11B genomic, mRNA and cDNA variants described herein.

Polynucleotides comprising a polymorphic gene variant or fragment of the invention may be useful for therapeutic purposes. For example, where a patient could benefit from expression, or increased expression, of a particular TNFRSF11B protein isoform, an expression vector encoding the isoform may be administered to the patient. The patient may be one who lacks the TNFRSF11B isogene encoding that isoform or may already have at least one copy of that isogene.

In other situations, it may be desirable to decrease or block expression of a particular TNFRSF11B isogene. Expression of a TNFRSF11B isogene may be turned off by transforming a targeted organ, tissue or cell population with an expression vector that expresses high levels of untranslatable mRNA or antisense RNA for the isogene or fragment thereof. Alternatively, oligonucleotides directed against the regulatory regions (e.g., promoter, introns, enhancers, 3' untranslated region) of the isogene may block transcription. Oligonucleotides targeting the transcription initiation site, e.g., between positions -10 and +10 from the start site are preferred. Similarly, inhibition of transcription can be achieved using oligonucleotides that base-pair with region(s) of the isogene DNA to form triplex DNA (see e.g., Gee et al. in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, N.Y., 1994). Antisense oligonucleotides may also be designed to block translation of TNFRSF11B mRNA transcribed from a particular isogene. It is also contemplated that ribozymes may be designed that can catalyze the specific cleavage of TNFRSF11B mRNA transcribed from a particular isogene.

The untranslated mRNA, antisense RNA or antisense oligonucleotides may be delivered to a target cell or tissue by expression from a vector introduced into the cell or tissue *in vivo* or *ex vivo*. Alternatively, such molecules may be formulated as a pharmaceutical composition for administration to the patient. Oligoribonucleotides and/or oligodeoxynucleotides intended for use as antisense oligonucleotides may be modified to increase stability and half-life. Possible modifications include, but are not limited to phosphorothioate or 2' O-methyl linkages, and the inclusion of nontraditional bases such as inosine and queosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytosine, guanine, thymine, and uracil which are not as easily recognized by endogenous nucleases.

The invention also provides an isolated polypeptide comprising a polymorphic variant of (a) the reference TNFRSF11B amino acid sequence shown in Figure 3 or (b) a fragment of this reference sequence. The location of a variant amino acid in a TNFRSF11B polypeptide or fragment of the invention is preferably identified by aligning its sequence against SEQ ID NO:3 (Fig. 3). A TNFRSF11B protein variant of the invention comprises an amino acid sequence identical to SEQ ID NO:3 for those regions of SEQ ID NO:3 that are encoded by examined portions of the TNFRSF11B gene (as described in the Examples below), except for having one or more variant amino acids selected from the group consisting of methionine at a position corresponding to amino acid position 240 and methionine at a position corresponding to amino acid position 281, and may also comprise an additional variant amino acid of

asparagine at a position corresponding to amino acid position 3. Thus, a TNFRSF11B fragment of the invention, also referred to herein as a TNFRSF11B peptide variant, is any fragment of a TNFRSF11B protein variant that contains one or more of the amino acid variations shown in Table 2. The invention specifically excludes amino acid sequences identical to those previously identified for TNFRSF11B, including SEQ ID NO:3, and previously described fragments thereof. TNFRSF11B protein variants included within the invention comprise all amino acid sequences based on SEQ ID NO:3 and having the combination of amino acid variations described in Table 2 below. In preferred embodiments, a TNFRSF11B protein variant of the invention is encoded by an isogene defined by one of the observed haplotypes, 3, 7, 8, 12-14, 18 and 22, shown in Table 5.

Table 2. Novel Polymorphic Variants of TNFRSF11B

Polymorphic Variant Number	Amino Acid Position and Identities		
	3	240	281
1	K	I	M
2	K	M	V
3	K	M	M
4	N	I	V
5	N	I	M
6	N	M	V
7	N	M	M

A TNFRSF11B peptide variant of the invention is at least 6 amino acids in length and is preferably any number between 6 and 30 amino acids long, more preferably between 10 and 25, and most preferably between 15 and 20 amino acids long. Such TNFRSF11B peptide variants may be useful as antigens to generate antibodies specific for one of the above TNFRSF11B isoforms. In addition, the TNFRSF11B peptide variants may be useful in drug screening assays.

A TNFRSF11B variant protein or peptide of the invention may be prepared by chemical synthesis or by expressing an appropriate variant TNFRSF11B genomic or cDNA sequence described above. Alternatively, the TNFRSF11B protein variant may be isolated from a biological sample of an individual having a TNFRSF11B isogene which encodes the variant protein. Where the sample contains two different TNFRSF11B isoforms (i.e., the individual has different TNFRSF11B isogenes), a particular TNFRSF11B isoform of the invention can be isolated by immunoaffinity chromatography using an antibody which specifically binds to that particular TNFRSF11B isoform but does not bind to the other TNFRSF11B isoform.

The expressed or isolated TNFRSF11B protein or peptide may be detected by methods known in the art, including Coomassie blue staining, silver staining, and Western blot analysis using antibodies

specific for the isoform of the TNFRSF11B protein or peptide as discussed further below. TNFRSF11B variant proteins and peptides can be purified by standard protein purification procedures known in the art, including differential precipitation, molecular sieve chromatography, ion-exchange chromatography, isoelectric focusing, gel electrophoresis, affinity and immunoaffinity chromatography and the like.

5 (Ausubel et. al., 1987, In Current Protocols in Molecular Biology John Wiley and Sons, New York, New York). In the case of immunoaffinity chromatography, antibodies specific for a particular polymorphic variant may be used.

10 A polymorphic variant TNFRSF11B gene of the invention may also be fused in frame with a heterologous sequence to encode a chimeric TNFRSF11B protein. The non-TNFRSF11B portion of the chimeric protein may be recognized by a commercially available antibody. In addition, the chimeric protein may also be engineered to contain a cleavage site located between the TNFRSF11B and non-TNFRSF11B portions so that the TNFRSF11B protein may be cleaved and purified away from the non-TNFRSF11B portion.

15 An additional embodiment of the invention relates to using a novel TNFRSF11B protein isoform, or a fragment thereof, in any of a variety of drug screening assays. Such screening assays may be performed to identify agents that bind specifically to all known TNFRSF11B protein isoforms or to only a subset of one or more of these isoforms. The agents may be from chemical compound libraries, peptide libraries and the like. The TNFRSF11B protein or peptide variant may be free in solution or affixed to a solid support. In one embodiment, high throughput screening of compounds for binding to a TNFRSF11B  
20 variant may be accomplished using the method described in PCT application WO84/03565, in which large numbers of test compounds are synthesized on a solid substrate, such as plastic pins or some other surface, contacted with the TNFRSF11B protein(s) of interest and then washed. Bound TNFRSF11B protein(s) are then detected using methods well-known in the art.

25 In another embodiment, a novel TNFRSF11B protein isoform may be used in assays to measure the binding affinities of one or more candidate drugs targeting the TNFRSF11B protein.

In yet another embodiment, when a particular TNFRSF11B haplotype or group of TNFRSF11B haplotypes encodes a TNFRSF11B protein variant with an amino acid sequence distinct from that of TNFRSF11B protein isoforms encoded by other TNFRSF11B haplotypes, then detection of that particular TNFRSF11B haplotype or group of TNFRSF11B haplotypes may be accomplished by detecting  
30 expression of the encoded TNFRSF11B protein variant using any of the methods described herein or otherwise commonly known to the skilled artisan.

In another embodiment, the invention provides antibodies specific for and immunoreactive with one or more of the novel TNFRSF11B protein or peptide variants described herein. The antibodies may be either monoclonal or polyclonal in origin. The TNFRSF11B protein or peptide variant used to generate the

antibodies may be from natural or recombinant sources (*in vitro* or *in vivo*) or produced by chemical synthesis or semi-synthetic synthesis using synthesis techniques known in the art. If the TNFRSF11B protein or peptide variant is of insufficient size to be antigenic, it may be concatenated or conjugated, complexed, or otherwise covalently linked to a carrier molecule to enhance the antigenicity of the peptide.

5 Examples of carrier molecules, include, but are not limited to, albumins (e.g., human, bovine, fish, ovine), and keyhole limpet hemocyanin (Basic and Clinical Immunology, 1991, Eds. D.P. Stites, and A.I. Terr, Appleton and Lange, Norwalk Connecticut, San Mateo, California).

In one embodiment, an antibody specifically immunoreactive with one of the novel protein or peptide variants described herein is administered to an individual to neutralize activity of the TNFRSF11B isoform expressed by that individual. The antibody may be formulated as a pharmaceutical composition which includes a pharmaceutically acceptable carrier.

10

Antibodies specific for and immunoreactive with one of the novel protein isoforms described herein may be used to immunoprecipitate the TNFRSF11B protein variant from solution as well as react with TNFRSF11B protein isoforms on Western or immunoblots of polyacrylamide gels on membrane supports or substrates. In another preferred embodiment, the antibodies will detect TNFRSF11B protein isoforms in paraffin or frozen tissue sections, or in cells which have been fixed or unfixed and prepared on slides, coverslips, or the like, for use in immunocytochemical, immunohistochemical, and immunofluorescence techniques.

15

In another embodiment, an antibody specifically immunoreactive with one of the novel TNFRSF11B protein variants described herein is used in immunoassays to detect this variant in biological samples. In this method, an antibody of the present invention is contacted with a biological sample and the formation of a complex between the TNFRSF11B protein variant and the antibody is detected. As described, suitable immunoassays include radioimmunoassay, Western blot assay, immunofluorescent assay, enzyme linked immunoassay (ELISA), chemiluminescent assay, immunohistochemical assay, immunocytochemical assay, and the like (see, e.g., Principles and Practice of Immunoassay, 1991, Eds. Christopher P. Price and David J. Neoman, Stockton Press, New York, New York; Current Protocols in Molecular Biology, 1987, Eds. Ausubel et al., John Wiley and Sons, New York, New York). Standard techniques known in the art for ELISA are described in Methods in Immunodiagnosis, 2nd Ed., Eds. Rose and Bigazzi, John Wiley and Sons, New York 1980; and Campbell et al., 1984, Methods in Immunology, W.A. Benjamin, Inc.). Such assays may be direct, indirect, competitive, or noncompetitive as described in the art (see, e.g., Principles and Practice of Immunoassay, 1991, Eds. Christopher P. Price and David J. Neoman, Stockton Pres, NY, NY; and Oellrich, M., 1984, J. Clin. Chem. Clin. Biochem., 22:895-904). Proteins may be isolated from test specimens and biological samples by conventional methods, as described in Current Protocols in Molecular Biology, supra.

20

25

30

Exemplary antibody molecules for use in the detection and therapy methods of the present invention are intact immunoglobulin molecules, substantially intact immunoglobulin molecules, or those portions of immunoglobulin molecules that contain the antigen binding site. Polyclonal or monoclonal antibodies may be produced by methods conventionally known in the art (e.g., Kohler and Milstein, 1975, Nature, 256:495-497; Campbell Monoclonal Antibody Technology, the Production and Characterization of Rodent and Human Hybridomas, 1985, In: Laboratory Techniques in Biochemistry and Molecular Biology, Eds. Burdon et al., Volume 13, Elsevier Science Publishers, Amsterdam). The antibodies or antigen binding fragments thereof may also be produced by genetic engineering. The technology for expression of both heavy and light chain genes in *E. coli* is the subject of PCT patent applications, publication number WO 901443 and WO 9014424 and in Huse et al., 1989, *Science*, 246:1275-1281. The antibodies may also be humanized (e.g., Queen, C. et al. 1989 *Proc. Natl. Acad. Sci. USA* 86:10029).

Effect(s) of the polymorphisms identified herein on expression of TNFRSF11B may be investigated by various means known in the art, such as by *in vitro* translation of mRNA transcripts of the TNFRSF11B gene, cDNA or fragment thereof, or by preparing recombinant cells and/or nonhuman recombinant organisms, preferably recombinant animals, containing a polymorphic variant of the TNFRSF11B gene. As used herein, "expression" includes but is not limited to one or more of the following: transcription of the gene into precursor mRNA; splicing and other processing of the precursor mRNA to produce mature mRNA; mRNA stability; translation of the mature mRNA(s) into TNFRSF11B protein(s) (including effects of polymorphisms on codon usage and tRNA availability); and glycosylation and/or other modifications of the translation product, if required for proper expression and function.

To prepare a recombinant cell of the invention, the desired TNFRSF11B isogene, cDNA or coding sequence may be introduced into the cell in a vector such that the isogene, cDNA or coding sequence remains extrachromosomal. In such a situation, the gene will be expressed by the cell from the extrachromosomal location. In a preferred embodiment, the TNFRSF11B isogene, cDNA or coding sequence is introduced into a cell in such a way that it recombines with the endogenous TNFRSF11B gene present in the cell. Such recombination requires the occurrence of a double recombination event, thereby resulting in the desired TNFRSF11B gene polymorphism. Vectors for the introduction of genes both for recombination and for extrachromosomal maintenance are known in the art, and any suitable vector or vector construct may be used in the invention. Methods such as electroporation, particle bombardment, calcium phosphate co-precipitation and viral transduction for introducing DNA into cells are known in the art; therefore, the choice of method may lie with the competence and preference of the skilled practitioner. Examples of cells into which the TNFRSF11B isogene, cDNA or coding sequence may be introduced include, but are not limited to, continuous culture cells, such as COS, CHO, NIH/3T3, and primary or culture cells of the relevant tissue type, i.e., they express the TNFRSF11B isogene, cDNA or coding

sequence. Such recombinant cells can be used to compare the biological activities of the different protein variants.

Recombinant nonhuman organisms, i.e., transgenic animals, expressing a variant TNFRSF11B gene, cDNA or coding sequence are prepared using standard procedures known in the art. Preferably, a construct comprising the variant gene, cDNA or coding sequence is introduced into a nonhuman animal or an ancestor of the animal at an embryonic stage, i.e., the one-cell stage, or generally not later than about the eight-cell stage. Transgenic animals carrying the constructs of the invention can be made by several methods known to those having skill in the art. One method involves transfecting into the embryo a retrovirus constructed to contain one or more insulator elements, a gene or genes (or cDNA or coding sequence) of interest, and other components known to those skilled in the art to provide a complete shuttle vector harboring the insulated gene(s) as a transgene, see e.g., U.S. Patent No. 5,610,053. Another method involves directly injecting a transgene into the embryo. A third method involves the use of embryonic stem cells. Examples of animals into which the TNFRSF11B isogene, cDNA or coding sequences may be introduced include, but are not limited to, mice, rats, other rodents, and nonhuman primates (see "The Introduction of Foreign Genes into Mice" and the cited references therein, In: Recombinant DNA, Eds. J.D. Watson, M. Gilman, J. Witkowski, and M. Zoller; W.H. Freeman and Company, New York, pages 254-272). Transgenic animals stably expressing a human TNFRSF11B isogene, cDNA or coding sequence and producing the encoded human TNFRSF11B protein can be used as biological models for studying diseases related to abnormal TNFRSF11B expression and/or activity, and for screening and assaying various candidate drugs, compounds, and treatment regimens to reduce the symptoms or effects of these diseases.

An additional embodiment of the invention relates to pharmaceutical compositions for treating disorders affected by expression or function of a novel TNFRSF11B isogene described herein. The pharmaceutical composition may comprise any of the following active ingredients: a polynucleotide comprising one of these novel TNFRSF11B isogenes (or cDNAs or coding sequences); an antisense oligonucleotide directed against one of the novel TNFRSF11B isogenes, a polynucleotide encoding such an antisense oligonucleotide, or another compound which inhibits expression of a novel TNFRSF11B isogene described herein. Preferably, the composition contains the active ingredient in a therapeutically effective amount. By therapeutically effective amount is meant that one or more of the symptoms relating to disorders affected by expression or function of a novel TNFRSF11B isogene is reduced and/or eliminated. The composition also comprises a pharmaceutically acceptable carrier, examples of which include, but are not limited to, saline, buffered saline, dextrose, and water. Those skilled in the art may employ a formulation most suitable for the active ingredient, whether it is a polynucleotide, oligonucleotide, protein, peptide or small molecule antagonist. The pharmaceutical composition may be

administered alone or in combination with at least one other agent, such as a stabilizing compound. Administration of the pharmaceutical composition may be by any number of routes including, but not limited to oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, intradermal, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal.

5 Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

For any composition, determination of the therapeutically effective dose of active ingredient and/or the appropriate route of administration is well within the capability of those skilled in the art. For example, the dose can be estimated initially either in cell culture assays or in animal models. The animal  
10 model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. The exact dosage will be determined by the practitioner, in light of factors relating to the patient requiring treatment, including but not limited to severity of the disease state, general health, age, weight and gender of the patient, diet, time and frequency of administration, other drugs being taken by the patient, and  
15 tolerance/response to the treatment.

Any or all analytical and mathematical operations involved in practicing the methods of the present invention may be implemented by a computer. In addition, the computer may execute a program that generates views (or screens) displayed on a display device and with which the user can interact to view and analyze large amounts of information relating to the TNFRSF11B gene and its genomic variation,  
20 including chromosome location, gene structure, and gene family, gene expression data, polymorphism data, genetic sequence data, and clinical data population data (e.g., data on ethnogeographic origin, clinical responses, genotypes, and haplotypes for one or more populations). The TNFRSF11B polymorphism data described herein may be stored as part of a relational database (e.g., an instance of an Oracle database or a set of ASCII flat files). These polymorphism data may be stored on the computer's hard drive or may, for  
25 example, be stored on a CD-ROM or on one or more other storage devices accessible by the computer. For example, the data may be stored on one or more databases in communication with the computer via a network.

Preferred embodiments of the invention are described in the following examples. Other  
embodiments within the scope of the claims herein will be apparent to one skilled in the art from  
30 consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples.

## EXAMPLES

The Examples herein are meant to exemplify the various aspects of carrying out the invention and are not intended to limit the scope of the invention in any way. The Examples do not include detailed descriptions for conventional methods employed, such as in the performance of genomic DNA isolation, PCR and sequencing procedures. Such methods are well-known to those skilled in the art and are described in numerous publications, for example, Sambrook, Fritsch, and Maniatis, "Molecular Cloning: A Laboratory Manual", 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory Press, USA, (1989).

### EXAMPLE 1

This example illustrates examination of various regions of the TNFRSF11B gene for polymorphic sites.

#### Amplification of Target Regions

The following target regions of the TNFRSF11B gene were amplified using PCR primer pairs. The primers used for each region are represented below by providing the nucleotide positions of their initial and final nucleotides, which correspond to positions in SEQ ID NO:1 (Figure 1).

#### PCR Primer Pairs

Fragment No.	Forward Primer	Reverse Primer	PCR Product
Fragment 1	427-446	complement of 1018-996	592 nt
Fragment 2	692-711	complement of 1437-1412	746 nt
Fragment 3	777-798	complement of 1217-1195	441 nt
Fragment 4	1604-1627	complement of 2208-2186	605 nt
Fragment 5	5748-5771	complement of 6485-6464	738 nt
Fragment 6	8035-8059	complement of 8653-8632	619 nt
Fragment 7	9942-9964	complement of 10628-10605	687 nt

These primer pairs were used in PCR reactions containing genomic DNA isolated from immortalized cell lines for each member of the Index Repository. The PCR reactions were carried out under the following conditions:

Reaction volume	= 10 $\mu$ l
10 x Advantage 2 Polymerase reaction buffer (Clontech)	= 1 $\mu$ l
100 ng of human genomic DNA	= 1 $\mu$ l
10 mM dNTP	= 0.4 $\mu$ l
Advantage 2 Polymerase enzyme mix (Clontech)	= 0.2 $\mu$ l
Forward Primer (10 $\mu$ M)	= 0.4 $\mu$ l
Reverse Primer (10 $\mu$ M)	= 0.4 $\mu$ l
Water	= 6.6 $\mu$ l

#### Amplification profile:

97°C - 2 min. 1 cycle

5      97°C - 15 sec.  
70°C - 45 sec.      }      10 cycles  
72°C - 45 sec.

10      97°C - 15 sec.  
64°C - 45 sec.      }      35 cycles  
72°C - 45 sec.

#### Sequencing of PCR Products

The PCR products were purified using a Whatman/Polyfritronics 100 µl 384 well unfilter plate essentially according to the manufacturers protocol. The purified DNA was eluted in 50 µl of distilled water. Sequencing reactions were set up using Applied Biosystems Big Dye Terminator chemistry essentially according to the manufacturers protocol. The purified PCR products were sequenced in both directions using the primer sets described previously or those represented below by the nucleotide positions of their initial and final nucleotides, which correspond to positions in SEQ ID NO:1 (Figure 1).

Reaction products were purified by isopropanol precipitation, and run on an Applied Biosystems 3700 DNA Analyzer.

#### Sequencing Primer Pairs

	Fragment No.	Forward Primer	Reverse Primer
25	Fragment 1	452-471	complement of 990-972
	Fragment 2	758-778	complement of 1285-1264
	Fragment 3	811-830	complement of 1181-1162
	Fragment 4	1660-1679	complement of 2180-2161
	Fragment 5	5823-5842	complement of 6352-6331
30	Fragment 6	8067-8086	complement of 8578-8557
	Fragment 7	10029-10050	complement of 10548-10529

#### Analysis of Sequences for Polymorphic Sites

Sequence information for a minimum of 80 humans was analyzed for the presence of polymorphisms using the Polyphred program (Nickerson et al., *Nucleic Acids Res.* 14:2745-2751, 1997). The presence of a polymorphism was confirmed on both strands. The polymorphisms and their locations in the TNFRSF11B reference genomic sequence (SEQ ID NO:1) are listed in Table 3 below.

Table 3. Polymorphic Sites Identified in the TNFRSF11B Gene

	Polymorphic Site Number	PolyId(a)	Nucleotide Position	Reference Allele	Variant Allele	CDS Variant Position	AA Variant
5	PS1	3827227	504	G	T		
	PS2	3827231	717	C	T		
	PS3	3827233	744	G	T		
	PS4	3827235	778	T	C		
	PS5(R)	3827239	1009	G	C	9	K3N
10	PS6	3827243	1045	C	T		
	PS7	3827245	1122	G	A		
	PS8	3827284	1218	C	A		
	PS9	3827437	2014	C	T		
	PS10	3827444	2177	T	C		
15	PS11	3827446	5906	T	C		
	PS12	3827451	6010	C	T		
	PS13	3827457	8110	G	A		
	PS14	3827459	8333	C	T	699	N233N
	PS15	3827461	8354	A	G	720	I240M
20	PS16	3827463	8402	A	G	768	L256L
	PS17	3827465	8459	A	C		
	PS18	3827467	10203	G	A	841	V281M
	PS19	3827469	10512	T	C	1150	L384L

- 25 (a) PolyId is a unique identifier assigned to each PS by Genaissance Pharmaceuticals, Inc.  
(R) Reported previously.

#### EXAMPLE 2

30 This example illustrates analysis of the TNFRSF11B polymorphisms identified in the Index Repository for human genotypes and haplotypes.

The different genotypes containing these polymorphisms that were observed in unrelated members of the reference population are shown in Table 4 below, with the haplotype pair indicating the combination of haplotypes determined for the individual using the haplotype derivation protocol described below. In  
35 Table 4, homozygous positions are indicated by one nucleotide and heterozygous positions are indicated by two nucleotides. Missing nucleotides in any given genotype in Table 4 were inferred based on linkage disequilibrium and/or Mendelian inheritance.

Table 4 (Part 1). Genotypes and Haplotype Pairs Observed for TNFRSF11B Gene

Genotype Number	HAP Pair	Polymorphic Sites									
		PS1	PS2	PS3	PS4	PS5	PS6	PS7	PS8	PS9	PS10
5	1 1	G	C	G	C	C	C	G	C	C	T
	15 15	G	C	G	T	G	C	G	A	C	T
	19 19	G	C	G	T	G	C	G	C	C	T
	6 6	G	C	G	C	G	T	G	C	C	T
	12 12	G	C	G	T	C	C	G	C	T	C
10	19 16	G	C	G	T	G	C	G	C	C	T
	10 2	G	C	G	T/C	C	C	G	C	C	T
	1 5	G	C	G	C	C/G	C/T	G	C	C	T
	10 14	G	C	G	T	C/G	C	G	C/A	C	T
	19 14	G	C	G	T	G	C	G	C/A	C	T
15	19 13	G	C	G	T	G	C	G	C/A	C	T
	15 12	G	C	G	T	G/C	C	G	A/C	C/T	T/C
	1 21	G	C/T	G	C	C	C	G	C	C	T
	19 11	G	C	G	T	G/C	C	G	C	C	T
	15 6	G	C	G	T/C	G	C/T	G	A/C	C	T
20	1 6	G	C	G	C	C/G	C/T	G	C	C	T
	1 2	G	C	G	C	C	C	G	C	C	T
	19 4	G	C	G	T/C	G	C/T	G	C	C	T
	19 3	G	C	G	T/C	G	C/T	G/A	C	C	T
	19 9	G	C	G	T	G/C	C	G	C/A	C	T
25	15 3	G	C	G	T/C	G	C/T	G/A	A/C	C	T
	19 20	G	C	G/T	T	G	C	G	C	C	T
	19 7	G	C	G	T/C	G	C/T	G	C	C/T	T/C
	19 18	G	C	G	T	G	C	G	C	C	T
	22 17	T/G	C	G	T	G	C	G	C	C	T
30	19 12	G	C	G	T	G/C	C	G	C	C/T	T/C
	1 12	G	C	G	C/T	C	C	G	C	C/T	T/C
	19 8	G	C	G	T	G/C	C	G	C/A	C	T
	15 10	G	C	G	T	G/C	C	G	A/C	C	T
	19 15	G	C	G	T	G	C	G	C/A	C	T
35	19 10	G	C	G	T	G/C	C	G	C	C	T
	18 16	G	C	G	T	G	C	G	C	C	T
	3 14	G	C	G	C/T	G	T/C	A/G	C/A	C	T

Table 4 (Part 2). Genotypes and Haplotype Pairs Observed for TNFRSF11B Gene

5	Genotype		Polymorphic Sites								
	Number	HAP Pair	PS11	PS12	PS13	PS14	PS15	PS16	PS17	PS18	PS19
10	1	1 1	C	C	G	C	A	A	A	G	T
	2	15 15	T	C	G	C	A	A	A	G	T
	3	19 19	T	C	G	C	A	A	A	G	T
	4	6 6	T	T	G	C	A	A	A	G	T
	5	12 12	T	C	G	C	A	G	C	G	T
	6	19 16	T/C	C	G	C	A	A	A	G	T
	7	10 2	C/T	C	G	C	A	A	A	G	T
	8	1 5	C/T	C	G	C	A	A	A	G	T
	9	10 14	C/T	C	G	C	A	A	A	G	T/C
15	10	19 14	T	C	G	C	A	A	A	G	T/C
	11	19 13	T/C	C	G/A	C	A	A	A	G	T/C
	12	15 12	T	C	G	C	A	A/G	A/C	G	T
	13	1 21	C	C	G	C	A	A	A	G	T
	14	19 11	T	C	G	C	A	A	A	G/A	T
20	15	15 6	T	C/T	G	C	A	A	A	G	T
	16	1 6	C/T	C/T	G	C	A	A	A	G	T
	17	1 2	C/T	C	G	C	A	A	A	G	T
	18	19 4	T/C	C	G	C	A	A	A	G	T
	19	19 3	T	C	G	C	A	A	A	G	T/C
25	20	19 9	T	C	G	C	A	A	A	G	T
	21	15 3	T	C	G	C	A	A	A	G	T/C
	22	19 20	T	C	G	C	A	A	A	G	T
	23	19 7	T	C/T	G	C/T	A	A/G	A	G	T
	24	19 18	T	C	G	C	A	A	A	G	T/C
30	25	22 17	T	C	A	C	A	A	A	G	C/T
	26	19 12	T	C	G	C	A	A/G	A/C	G	T
	27	1 12	C/T	C	G	C	A	A/G	A/C	G	T
	28	19 8	T/C	C	G	C	A/G	A	A	G	T
	29	15 10	T/C	C	G	C	A	A	A	G	T
35	30	19 15	T	C	G	C	A	A	A	G	T
	31	19 10	T/C	C	G	C	A	A	A	G	T
	32	18 16	T/C	C	G	C	A	A	A	G	C/T
	33	3 14	T	C	G	C	A	A	A	G	C

40 The haplotype pairs shown in Table 4 were estimated from the unphased genotypes using a computer-implemented extension of Clark's algorithm (Clark, A.G. 1990 *Mol Bio Evol* 7, 111-122) for assigning haplotypes to unrelated individuals in a population sample, as described in PCT/US01/12831, filed April 18, 2001. In this method, haplotypes are assigned directly from individuals who are homozygous at all sites or heterozygous at no more than one of the variable sites. This list of haplotypes is

45 then used to deconvolute the unphased genotypes in the remaining (multiply heterozygous) individuals. In the present analysis, the list of haplotypes was augmented with haplotypes obtained from two families (one three-generation Caucasian family and one two-generation African-American family).

By following this protocol, it was determined that the Index Repository examined herein and, by extension, the general population contains the 22 human TNFRSF11B haplotypes shown in Table 5 below.

A TNFRSF11B isogene defined by a full-haplotype shown in Table 5 below comprises the regions of the SEQ ID NOS indicated in Table 5, with their corresponding set of polymorphic locations and identities, which are also set forth in Table 5.

Table 5(Part 1). Haplotypes of the TNFRSF11B gene.

Regions	PS	PS	Haplotype Number(d)									
Examined(a)	No.(b)	Position(c)	1	2	3	4	5	6	7	8	9	10
10 427-1437	1	504/30	G	G	G	G	G	G	G	G	G	G
427-1437	2	717/150	C	C	C	C	C	C	C	C	C	C
427-1437	3	744/270	G	G	G	G	G	G	G	G	G	G
427-1437	4	778/390	C	C	C	C	C	C	C	T	T	T
427-1437	5	1009/510	C	C	G	G	G	G	G	C	C	C
15 427-1437	6	1045/630	C	C	T	T	T	T	T	C	C	C
427-1437	7	1122/750	G	G	A	G	G	G	G	G	G	G
427-1437	8	1218/870	C	C	C	C	C	C	C	A	A	C
1604-2208	9	2014/990	C	C	C	C	C	C	T	C	C	C
1604-2208	10	2177/1110	T	T	T	T	T	T	C	T	T	T
20 5748-6485	11	5906/1230	C	T	T	C	T	T	T	C	T	C
5748-6485	12	6010/1350	C	C	C	C	C	T	T	C	C	C
8035-8653	13	8110/1470	G	G	G	G	G	G	G	G	G	G
8035-8653	14	8333/1590	C	C	C	C	C	C	T	C	C	C
8035-8653	15	8354/1710	A	A	A	A	A	A	A	G	A	A
25 8035-8653	16	8402/1830	A	A	A	A	A	A	G	A	A	A
8035-8653	17	8459/1950	A	A	A	A	A	A	A	A	A	A
9942-10628	18	10203/2070	G	G	G	G	G	G	G	G	G	G
9942-10628	19	10512/2190	T	T	C	T	T	T	T	T	T	T

Table 5(Part 2). Haplotypes of the TNFRSF11B gene.

Regions	PS	PS	Haplotype Number(d)									
Examined(a)	No.(b)	Position(c)	11	12	13	14	15	16	17	18	19	20
5	427-1437	1	504/30	G	G	G	G	G	G	G	G	G
	427-1437	2	717/150	C	C	C	C	C	C	C	C	C
	427-1437	3	744/270	G	G	G	G	G	G	G	G	T
	427-1437	4	778/390	T	T	T	T	T	T	T	T	T
	427-1437	5	1009/510	C	C	G	G	G	G	G	G	G
	427-1437	6	1045/630	C	C	C	C	C	C	C	C	C
10	427-1437	7	1122/750	G	G	G	G	G	G	G	G	G
	427-1437	8	1218/870	C	C	A	A	A	C	C	C	C
	1604-2208	9	2014/990	C	T	C	C	C	C	C	C	C
	1604-2208	10	2177/1110	T	C	T	T	T	T	T	T	T
	5748-6485	11	5906/1230	T	T	C	T	T	C	T	T	T
15	5748-6485	12	6010/1350	C	C	C	C	C	C	C	C	C
	8035-8653	13	8110/1470	G	G	A	G	G	G	A	G	G
	8035-8653	14	8333/1590	C	C	C	C	C	C	C	C	C
	8035-8653	15	8354/1710	A	A	A	A	A	A	A	A	A
	8035-8653	16	8402/1830	A	G	A	A	A	A	A	A	A
20	8035-8653	17	8459/1950	A	C	A	A	A	A	A	A	A
	9942-10628	18	10203/2070	A	G	G	G	G	G	G	G	G
	9942-10628	19	10512/2190	T	T	C	C	T	T	C	T	T

Table 5(Part 3). Haplotypes of the TNFRSF11B gene.

25	Regions	PS	PS	Haplotype Number(d)	
	Examined(a)	No.(b)	Position(c)	21	22
30	427-1437	1	504/30	G	T
	427-1437	2	717/150	T	C
	427-1437	3	744/270	G	G
	427-1437	4	778/390	C	T
	427-1437	5	1009/510	C	G
	427-1437	6	1045/630	C	C
	427-1437	7	1122/750	G	G
	427-1437	8	1218/870	C	C
35	1604-2208	9	2014/990	C	C
	1604-2208	10	2177/1110	T	T
	5748-6485	11	5906/1230	C	T
	5748-6485	12	6010/1350	C	C
40	8035-8653	13	8110/1470	G	A
	8035-8653	14	8333/1590	C	C
	8035-8653	15	8354/1710	A	A
	8035-8653	16	8402/1830	A	A
	8035-8653	17	8459/1950	A	A
	9942-10628	18	10203/2070	G	G
	9942-10628	19	10512/2190	T	C

(a) Region examined represents the nucleotide positions defining the start and stop positions within SEQ ID NO:1 of the regions sequenced;

(b) PS = polymorphic site;

- (c) Position of PS within the indicated SEQ ID NO, with the 1<sup>st</sup> position number referring to SEQ ID NO:1 and the 2<sup>nd</sup> position number referring to SEQ ID NO:94, a modified version of SEQ ID NO:1 that comprises the context sequence of each polymorphic site, PS1-PS19, to facilitate electronic searching of the haplotypes;
- 5 (d) Alleles for TNFRSF11B haplotypes are presented 5' to 3' in each column.

SEQ ID NO:1 refers to Figure 1, with the two alternative allelic variants of each polymorphic site indicated by the appropriate nucleotide symbol. SEQ ID NO:94 is a modified version of SEQ ID NO:1 that shows the context sequence of each of PS1-PS19 in a uniform format to facilitate electronic searching of the TNFRSF11B haplotypes. For each polymorphic site, SEQ ID NO:94 contains a block of 10 60 bases of the nucleotide sequence encompassing the centrally-located polymorphic site at the 30<sup>th</sup> position, followed by 60 bases of unspecified sequence to represent that each polymorphic site is separated by genomic sequence whose composition is defined elsewhere herein.

Table 6 below shows the percent of chromosomes characterized by a given TNFRSF11B 15 haplotype for all unrelated individuals in the Index Repository for which haplotype data was obtained. The percent of these unrelated individuals who have a given TNFRSF11B haplotype pair is shown in Table 7. In Tables 6 and 7, the "Total" column shows this frequency data for all of these unrelated individuals, while the other columns show the frequency data for these unrelated individuals categorized according to their self-identified ethnogeographic origin. Abbreviations used in Tables 6 and 7 are AF = African 20 Descent, AS = Asian, CA = Caucasian, HL = Hispanic-Latino, and AM = Native American.

10044674-010902

Table 6. Frequency of Observed TNFRSF11B Haplotypes In Unrelated Individuals

	HAP No.	HAP ID	Total	CA	AF	AS	HL	AM
5	1	37738726	13.41	19.05	0.0	0.0	30.56	50.0
	2	37738810	1.22	2.38	0.0	0.0	0.0	16.67
	3	37738753	3.66	0.0	15.0	0.0	0.0	0.0
	4	37738818	0.61	2.38	0.0	0.0	0.0	0.0
	5	37738904	0.61	0.0	0.0	0.0	0.0	16.67
10	6	37738747	4.88	7.14	7.5	0.0	5.56	0.0
	7	37738838	0.61	0.0	2.5	0.0	0.0	0.0
	8	37738876	0.61	0.0	0.0	2.5	0.0	0.0
	9	37738891	0.61	0.0	2.5	0.0	0.0	0.0
	10	37738714	14.02	16.67	5.0	17.5	16.67	16.67
15	11	37738825	0.61	2.38	0.0	0.0	0.0	0.0
	12	37738740	5.49	9.52	0.0	2.5	11.11	0.0
	13	37738853	0.61	0.0	2.5	0.0	0.0	0.0
	14	37738768	2.44	0.0	10.0	0.0	0.0	0.0
	15	37738707	14.63	16.67	10.0	15.0	19.44	0.0
20	16	37738797	1.22	0.0	2.5	0.0	2.78	0.0
	17	37738911	0.61	0.0	2.5	0.0	0.0	0.0
	18	37738781	1.22	0.0	5.0	0.0	0.0	0.0
	19	37738685	31.1	21.43	30.0	62.5	13.89	0.0
	20	37738882	0.61	0.0	2.5	0.0	0.0	0.0
25	21	37738862	0.61	2.38	0.0	0.0	0.0	0.0
	22	37738848	0.61	0.0	2.5	0.0	0.0	0.0

20044574-010902

Table 7. Frequency of Observed TNFRSF11B Haplotype Pairs In Unrelated Individuals

	HAP1	HAP2	Total	CA	AF	AS	HL	AM
5	1	1	7.32	9.52	0.0	0.0	16.67	33.33
	15	15	2.44	4.76	0.0	0.0	5.56	0.0
	19	19	10.98	4.76	10.0	30.0	0.0	0.0
	6	6	2.44	4.76	5.0	0.0	0.0	0.0
10	12	12	1.22	4.76	0.0	0.0	0.0	0.0
	19	16	1.22	0.0	0.0	0.0	5.56	0.0
	10	2	1.22	0.0	0.0	0.0	0.0	33.33
	1	5	1.22	0.0	0.0	0.0	0.0	33.33
	10	14	1.22	0.0	5.0	0.0	0.0	0.0
	19	14	1.22	0.0	5.0	0.0	0.0	0.0
15	19	13	1.22	0.0	5.0	0.0	0.0	0.0
	15	12	1.22	0.0	0.0	0.0	5.56	0.0
	1	21	1.22	4.76	0.0	0.0	0.0	0.0
	19	11	1.22	4.76	0.0	0.0	0.0	0.0
	15	6	1.22	0.0	5.0	0.0	0.0	0.0
20	1	6	3.66	4.76	0.0	0.0	11.11	0.0
	1	2	1.22	4.76	0.0	0.0	0.0	0.0
	19	4	1.22	4.76	0.0	0.0	0.0	0.0
	19	3	2.44	0.0	10.0	0.0	0.0	0.0
	19	9	1.22	0.0	5.0	0.0	0.0	0.0
	15	3	2.44	0.0	10.0	0.0	0.0	0.0
25	19	20	1.22	0.0	5.0	0.0	0.0	0.0
	19	7	1.22	0.0	5.0	0.0	0.0	0.0
	19	18	1.22	0.0	5.0	0.0	0.0	0.0
	22	17	1.22	0.0	5.0	0.0	0.0	0.0
	19	12	2.44	4.76	0.0	5.0	0.0	0.0
30	1	12	4.88	4.76	0.0	0.0	16.67	0.0
	19	8	1.22	0.0	0.0	5.0	0.0	0.0
	15	10	10.98	19.05	5.0	5.0	16.67	0.0
	19	15	8.54	4.76	0.0	25.0	5.56	0.0
	19	10	14.63	14.29	0.0	30.0	16.67	0.0
35	18	16	1.22	0.0	5.0	0.0	0.0	0.0
	3	14	2.44	0.0	10.0	0.0	0.0	0.0

The size and composition of the Index Repository were chosen to represent the genetic diversity across and within four major population groups comprising the general United States population. For example, as described in Table 1 above, this repository contains approximately equal sample sizes of African-descent, Asian-American, European-American, and Hispanic-Latino population groups. Almost all individuals representing each group had all four grandparents with the same ethnogeographic background. The number of unrelated individuals in the Index Repository provides a sample size that is sufficient to detect SNPs and haplotypes that occur in the general population with high statistical certainty. For instance, a haplotype that occurs with a frequency of 5% in the general population has a probability higher than 99.9% of being observed in a sample of 80 individuals from the general population. Similarly,

10044574 " 010902  
a haplotype that occurs with a frequency of 10% in a specific population group has a 99% probability of being observed in a sample of 20 individuals from that population group. In addition, the size and composition of the Index Repository means that the relative frequencies determined therein for the haplotypes and haplotype pairs of the TNFRSF11B gene are likely to be similar to the relative frequencies of these TNFRSF11B haplotypes and haplotype pairs in the general U.S. population and in the four population groups represented in the Index Repository. The genetic diversity observed for the three Native Americans is presented because it is of scientific interest, but due to the small sample size it lacks statistical significance.

In view of the above, it will be seen that the several advantages of the invention are achieved and other advantageous results attained.

As various changes could be made in the above methods and compositions without departing from the scope of the invention, it is intended that all matter contained in the above description and shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

All references cited in this specification, including patents and patent applications, are hereby incorporated in their entirety by reference. The discussion of references herein is intended merely to summarize the assertions made by their authors and no admission is made that any reference constitutes prior art. Applicants reserve the right to challenge the accuracy and pertinency of the cited references.